

เอกสาร

- กราฟอัตราการรอดชีวิตของผู้ป่วยโรคมะเร็งปากมดลูก ช่วง พ.ศ. 2493 - 2496
โรงพยาบาลหญิง (Survivorship Curves : Cervical Cancer Women's Hospital)
- สถิติการคลอดบุตรช่วง พ.ศ. 2493 - 2500 โรงพยาบาลหญิง (Obstetrical Statistics
1951 - 1957)
- สถิติผู้ป่วยโรคหูดหงอนไก่ที่อวัยวะสืบพันธุ์ ช่วง พ.ศ. 2494 - 2499 โรงพยาบาลหญิง
(Condyloma Acuminata)
- สถิติยุติกรรม(การแท้งลูก)ช่วง พ.ศ. 2494 - 2499 โรงพยาบาลหญิง (Abortion)
- สถิติผู้ป่วยโรคมะเร็งในกระเพาะอาหาร ช่วง พ.ศ. 2494 - 2500 โรงพยาบาลหญิง
(Malignancy of The stomach)
- สถิติผู้ป่วยโรคมะเร็งในเด็กและผู้หญิง ช่วง พ.ศ. 2494 - 2503 โรงพยาบาลหญิงและ
เด็ก
- สถิติทารกคลอดก่อนกำหนด ช่วง พ.ศ. 2495 - 2499 โรงพยาบาลหญิง
- สรุปสถิติของแผนกสูตินารีเวช ช่วง พ.ศ. 2496 - 2499 โรงพยาบาลหญิง (Statistical
Summary of Obstetrical Department in Women's Hospital)
- สถิติการคลอดผิดปกติ แผนกสูตินารีเวช พ.ศ. 2497 โรงพยาบาลหญิง
- สถิติผู้ป่วยโรคเหน็บชา ช่วง พ.ศ. 2496 - 2499

พ.ศ. 2493 - 2501

เอกสารส่วนบุคคล ศาสตราจารย์นายแพทย์เสม พริ้งพวงแก้ว

สพ 1

๒/๑.๑ การเกิดรายการรูดบัญชีของเงินโดยทางเงินฝากธนาคาร (ชื่อเรื่อง)

โรงพยาบาลสมเด็จ ร่อง พ.ศ. ๒๔๙๓-๒๔๙๖

พ.ศ. ๒๔๙๓ - ๒๔๙๖

จำนวน ๕ แผ่น


แฟ้มที่ ๒

กล่องที่ ๑

SURVIVORSHIP CURVES : CERVICAL CANCER

WOMEN'S HOSPITAL

*All cases.
(Including Stage 0).*



All cases

- CASES SEEN WITH VIEW TO TREATMENT -

	1950		1951		1952		1953		R.S.Clinic		Other		Total	
	R.S. Clinic	Other	R.S. Clinic	Other	R.S. Clinic	Other	R.S. Clinic	Other	Total	%	Total	%	No.	%
Treated	44	-	63	2	43	18	41	15	191	94.1	35	97.2	226	94.6
Untreated	2	-	4	-	5	-	1	1	12	5.9	1	2.8	13	5.4
Total	46	-	67	2	48	18	42	16	203	100.0	36	100.0	239	100.0

- DISTRIBUTION OF STAGE GROUPS -

Stage	1950		1951		1952		1953		R.S.Clinic		Other		Total	
	R.S. Clinic	Other	R.S. Clinic	Other	R.S. Clinic	Other	R.S. Clinic	Other	Total	%	Total	%	Total	%
I	17	-	19	2	21	5	10	8	67	33.0	15	41.7	82	34.3
II	20	-	36	-	17	8	23	3	96	47.3	11	30.5	107	44.8
III	6	-	8	-	8	2	6	3	28	13.8	5	13.9	33	13.8
	3	-	4	-	1	2	3	2	11	5.4	4	11.1	15	6.3
	-	-	-	-	1	1	-	-	1	0.5	1	2.8	2	0.8
Total	46	-	67	2	48	18	42	16	203	100.0	36	100.0	239	100.0

- RESIDUAL GROWTH - (Cases treated by Radio-therapy and Surgery).

	Total	
	No.	%
Observations	70	100.0
Positive	19	27.1

- - - - -

- NODE METASTASES - (Cases in which lymphadenectomy was carried out).

	Total	
	No.	%
Observations	123	100.0
Positive	26	21.1

- - - - -

ALTERATION OF STAGING AFTER OPERATION

Stage after Operation	Stage prior to operation				
	I	II	III	IV	Total
I	<u>45</u>	5	-	-	50
II	7	<u>38</u>	2	-	47
III	4	15	<u>3</u>	-	22
IV	-	-	1	<u>5</u>	6
Total	56	58	6	5	125

Overstaged prior to operation - 7 i.e. 5.6%
Understaged prior to operation - 27 i.e. 21.6%
Stage shown to be correct at operation - 91 i.e. 72.8%

N.B. Not included in the above table:-

2 Cases staged prior to operation which most probably
were actually Stage I and Stage III. The evidence
available, however, is not conclusive.

- TYPE OF OPERATIONS -

Type of Operations	1950		1951		1952		1953		Total	
	R.S. Clinic	Other	R.S. Clinic	Other	R.S. Clinic	Other	R.S. Clinic	Other	R.S. Clinic	Other
Wertheim	16	-	28	1	22	9	27	7	93	17
Anterior Exenteration	1	-	1	-	-	1	3	-	5	1
Palliative Gilchrist Op ⁿ .	-	-	-	-	-	-	1	-	1	-
Total Hysterectomy & Removal Adnexa.	-	-	-	-	-	2	-	1	-	3
Ligation of Hypogastric Arteries.	-	-	-	-	1	-	-	-	1	-
Extra peritoneal Lymphadenectomy.	1	-	4	-	-	-	-	-	5	-
Total Hysterectomy, Removal Adnexa & Pelvic Lymphadenectomy.	-	-	-	1	-	-	-	-	-	1
Total operations	18	-	33	2	23	12	31	8	105	22

- TYPE OF TREATMENT -

	1950		1951		1952		1953		R.S. Clinic		Other		Total	
	R.S. Clinic	Other	R.S. Clinic	Other	R.S. Clinic	Other	R.S. Clinic	Other	Total	%	Total	%	No.	%
Not treated	2	-	4	-	5	-	1	1	12	5.9	1	2.8	13	5.4
Radiotherapy -														
Radical	22	-	24	-	17	5	8	3	71	35.1	8	22.2	79	33.0
Palliative	4	-	6	-	3	1	3	4	16	7.9	5	13.5	21	8.8
Surgery -														
Radical	6	-	23	2	6	8	1	-	36	17.7	10	27.8	46	19.2
Palliative	-	-	-	-	1	1	1	-	2	1.0	1	2.8	3	1.3
Surgery & Radiotherapy -														
Radical	10	-	10	-	16	3	27	8	63	31.0	11	30.5	74	31.0
Palliative	2	-	-	-	-	-	1	-	3	1.4	-	-	3	1.3
	46	-	67	2	48	18	42	16	203	100.0	36	100.0	239	100.0

N.B. Figures shown thus () are the Other clinics, others are the R.S. Clinic.

	Stage I		Stage II		Stage III		Stage IV		All Stages	
	Alive	Dead	Alive	Dead	Alive	Dead	Alive	Dead	Alive	Dead
Not treated	-	-	-	-	-	-	-	2	-	2
Radio-therapy -										
Radical	3	4	10	3	-	2	-	-	13	9
Palliative	-	-	-	1	-	3	-	-	-	4
Surgery -										
Radical	4	-	1	1	-	-	-	-	5	1
Palliative	-	-	-	-	-	-	-	-	-	-
Surgery & Radio-therapy -										
Radical	5	1	2	2	-	-	-	-	7	3
Palliative	-	-	-	-	-	1	-	1	-	2
Total	12	5	13	7	-	6	-	3	25	21

1951

	Stage I		Stage II		Stage III		Stage IV		All Stages	
	Alive	Dead	Alive	Dead	Alive	Dead	Alive	Dead	Alive	Dead
Not treated	-	-	-	-	-	2	-	2	-	4
Radio-therapy -										
Radical	5	2	7	7	1	2	-	-	13	11
Palliative	-	-	-	1	-	3	-	2	-	6
-	(1)5	(1)6	5	7	-	-	-	-	(1)10	(1)13
Radical	-	-	-	-	-	-	-	-	-	-
Palliative	-	-	-	-	-	-	-	-	-	-
Surgery & Radio-therapy -										
Radical	1	-	4	5	-	-	-	-	5	5
Palliative	-	-	-	-	-	-	-	-	-	-
Total	(1)11	(1)8	16	20	1	7	-	4	(1)28	(1)39

1952

	Stage I		Stage II		Stage III		Stage IV		All Stages	
	Alive	Dead	Alive	Dead	Alive	Dead	Alive	Dead	Alive	Dead
Not treated	-	-	-	-	-	4	-	-	-	4
Radio-therapy -										
Radical	(1)5	2	(1)6	(2)4	-	(1)2	-	-	(2)11	(2)6
Palliative	-	-	-	-	-	2	-	1	-	(1)3
Surgery -										
Radical	(3)4	1	(4)1	-	-	(1)-	-	(1)-	(7)5	(1)1
Palliative	-	-	-	-	-	1	-	(1)-	-	(1)1
Surgery & Radio-therapy -										
Radical	(1)8	1	(1)5	1	1	-	-	(1)-	(2)14	(1)2
Palliative	-	-	-	-	-	-	-	-	-	-
Total	(3)17	4	(6)12	(2)5	1	(2)7	-	(2)1	(11)30	(6)17

x Includes 1 not traced.

N.B.

2 unstaged cases are not included.
1 not treated - dead R.S. Clinic
1 Radical Radio-therapy - dead - other clinic.

Yucca latex
50 cc — 1000 ~~cc~~
plasma

Calcium chloride

3%, 80. 1.022

50 cc / 1000 ~~cc~~
plasma

Othman Plasma. — First layer
out of 2 blue
acc.

Fractionation — no difference in
middle of collection

Fraction 1 — fibrinogen
& antibodies plus
glob.

Fraction 2 — 0 gam. glob.
& antibodies.

③ 50 agglutinin

Fraction 3 — discarded.

Fraction 4 — albumin

0.28 glob.
discarded

Alcohol, as Precipitation

M. F. J. DEMPSTER.

Dr. J. J. GRAYDON

เอกสารส่วนบุคคล ศาสตราจารย์นายแพทย์เสม พริ้งพวงแก้ว

สพ 1

..... 2/1.2 สภามหาวิทยาลัยแพทยศาสตร์ 2493-2500 (ชื่อเรื่อง)
↑
โรงพยาบาลสมเด็จ

พ.ศ. 2493-2500

จำนวน 9 แผ่น

แฟ้มที่ 2

กล่องที่ 1

O.B. Statistics 1958

2500

Dr. A.	2500	
Abnormal	Abnormal	Abnormal
200000	729	36
200000	695	38
200000	834	52
200000	814	60
200000	829	55
200000	881	55
200000	868	54
200000	909	67
200000	951	68
200000	973	86
200000	997	76
200000	1002	69
Total	<u>10482</u>	<u>716</u>

Annual Report of Women's Hospital 1956

Obstetrical Statistic (Women's Hospital 1951 - 1956)


Total admission	33,752	cases
A. <u>Normal full term delivery</u>	30,412	"
<u>Abnormal delivery</u>	3,340	"
(Including cesarean S. Forcep ext. Breech extraction craneotomy & internal version)		
Dead babies (full term)	626	(1:54)

B. Premature

In 1951 we considered baby who had weight under 2500 gm. to be premature. After one year we have found that the babies which had weight 2500 gm to 2200 gm could alive, 90 percenta, so we reconsidered the 2200 gm to be the weight of premature baby.

Premature (Wt. under 2500 gm. 1951)	256	(1:54)
Premature wt. under 2200 gm. 1952 - 1956	1337	(1:22)
Dead babies	152	(1:10)

Diagram shows numbers of delivery 1951 - 1956



Multiple Pregnancy

Twins	318	(1:106)
Triplets	8	(1:4219)
Quadruplets	1	(1: 33752)
Simese Twins	2	(1: 16876)
(Thoracophagus)		

Presentation

O L A	(in 6000 cases)	64.6 %
O R A	(in 6000 cases)	34.1 %
Breech	(1 : 90)	0.9 %
O R P, O P P, O L P	(1:200)	0.53%
	Transverse (1:719)	0.14 %
(Brow 1 cases)	Face & Brow (1 : 1125)	0.09 %

An average weight of new born	(in 6000 cases)	3068.4 gm
" " length of " "	(in 4000 ")	49.36 cm.
" " " " umbilical cord	(in 4000 cases)	49.74)"
" " weight of Placenta	(in 4000 cases)	579.6 gm.

An average of external pelvic measurement

Interspinal diameter	23.6 cm.	(in 6000 cases)
Intercristal "	26.11 cm.	(")
External conjugde	18.03 cm.	(")

Blood Kahn positive	7.8 %
---------------------	-------

OPERATIVE DELIVERY

(A) Cesarean Section 764 cases = 1:44
= 2.2 %

Indication	Number
Prolong labour. (1 st stage, 2 nd stage rigid cervix	229
Placenta previa	170
General Contracted pelvis & Flat pelvis	64
Toxemia of Pregnancy (eclampsia c severe Freeclampsia)	81
Obstructed labour (abnormal presentation Hydrocephalus, overterm)	69
Twins	7
Threatened uteri rupture	24
Heart disease	6
Failed forcep	4
Eldery primigravida (31 - 41)	28
Abruptio placenta	19
No reports	6
Descensus cervicis	1
Epilepsy	2
Previous cesarean section (6 months - 1 year)	31
Intraligamentary pregnancy	1
Missed abortion	1
Condyloma of vulva	2
Foetal distress (due to cord round the neck)	3
Bleeding from venous plexus (of uterus)	1
Antepartum bleeding	1
Profuse bleeding (Hypofibrinogenemia)	1
Prolapse cord	2
Appendicial abscess (Provisional diag. Threatened uteri rupture)	1
Total	764

Cesarean section

Maternal Mortality 14 = 1:55 = 1.8 %
Foetal Mortality 47 = 1:17 = 6 %

(2) Cesarean section hysterectomy

Uteri rupture	28	
Placenta accreta	2	
Total	30	
Maternal Mortality	3	(Uteri rupture)
Foetal Mortality	28	

(3) Forcep extraction 605 cases

1:55 = 1.8 %

Indication	Number
Uteri inertia	532
Toxemia of Pregnancy (Eclampsia)	16
Elderly primigravidarum c uteri inertia	7
Abnormal rotation of Head (O P P, O L P)	12
Twins	2
Deep transverse arrest	6
Pendulous abdomen	2
Heart disease	4
Hydrops fetalis	1
Abruptio placentae	6
Previous cesarean section	2
Out let forcip	6
Pulm T.B, c uteri inertia	2
Prolapse cord	1
Threatened uteri ruptured	2
No reports	4
Total	605

Maternal Mortality (2) = 1: 300 = 0.3 %
Foetal Mortality 34 = 1: 19 = 5.0 %

(4) Internal version

67 cases

Indication	Number
Multiple pregnancy	32
Prolapse arm	5
Transverse presentation	17
Placenta previa	2
Prolapse cord	1
Total	67

Maternal Mortality

Foetal Mortality 15 = 1:2 = 25 %

(5) Craniotomy

17

cases

Indication	Number
Obstructed labour due to wrong management (private midwife)	1
Hydrocephalus	1
Prolapse cord & dead foetus	1
Multiple pregnancy & dead foetus	3
Dead baby & abnormal rotation of head	1
Dead foetus in utero	1
No reports	3
Total	17

Maternal Mortality

nil.

(6) Breech extraction

380 cases

incidence

= 1 : 90

= 0.9 %

Maternal Complications

	Number	incidence	%
Placenta previa	188	1:200	0.5
Post partum hemorrhage	302	1:112	0.9
Uteri rupture	28	1:1200	0.1
Abruptio placentae	41	1:820	0.1
Toxemia of pregnancy	204	1:111	0.6
Placenta accreta	2	1:15371	0.01
Hypofibrinogenemia	1	1:33752	0.002
Hydrops gravidarum	14	1:2332	0.04
Bicoenu uterus	2	1:15371	0.01
Insertio velamentosa	1	1:33752	0.002
Polyhydramnios	4	1:8438	0.02
Threatened uteri rupture	19	1:1776	0.05
Miscellaneous	224	1:150	0.7
Retained Placenta	79	1: 427	0.2

Anomaly of Babies

	Number
Polydactylism	19
Harelip	19
Harelip c cleft palate	13
Talipes valgus	7
Hydrocephalus (frontalmeningocele 1)	13
Umbilical hernia	4
Monster	79
Hepatomegaly	1
Bow leg	1
Hydrocele c undescend testis	3
Diaphragmatic hernia	1
Stiff of knee joint	1
Situs inversus	1
Osteogenesis imperfecta	1
Siamese Twins	2
Imperforate ani	4
Hernia	2
Phimosis	1
Congenital megacolon	1
Facial paralysis	1
Recto vaginal fistula	1
Inguinal hernia	1
Cystic hygroma	1
Meningocele	1

Mortality

Mother 51 = 1: 660 = 0.1 %

Cause of Death	Number
Uteri rupture	4
Tear cervix	1
Toxemia of Pregnancy	10
Thrombocytopenic purpura	1
Severe shock	12
Uremia (after cesarean section)	1
Post partum hemorrhage	11
Intestinal obstruction	1
Peritonitis (after cesarean section)	2
Hypofibrinogenemia	1
Uteri atony	4
Heart failure	3
Renal failure	1

Baby 626 = 1:54 = 1.9 %

Cause of Death	Number
Macerated	278
Still birth	92
Asphyxia	22
Monster	82
Abnormal delivery	111
Dead during post partum	41

B.E. 2500

สถิติต่าง ๆ ของกรมการแพทย์

โรงพยาบาลทั่วไปมีเตียง ๔๔๕๐ เตียง

โรงพยาบาลโรคจิตมีเตียง ๔๒๕๐ เตียง
รวม ๘๗๐๐ เตียง

จำนวนพลเมือง ๒๐ ล้าน อัตราเตียง • เตียง ต่อประชาชน ๑๐๐๐ คน

๒๐ ล้าน จึงควรมีเตียง ๒๐,๐๐๐ เตียง

ค่าใช้จ่ายของเตียงคนไข้ใน • ปี ค.ศ. ๒๔๘๕ = ๒๔๘๔ บาท

เอกสารส่วนบุคคล ศาสตราจารย์นายแพทย์เสม พริ้งพวงแก้ว

สภ 1

. 2/1.3 สติจิตตมนอนอนิกกิจฉะสันตสิริ (ชื่อเรื่อง)

ช่วง พ.ศ. 2494 - 2499 โรงพยาบาลศิริราช

พ.ศ. 2494 - 2499

จำนวน 15 แผ่น

แฟ้มที่ 2

กล่องที่ 1

2494 - 2498.

30 N 0 N Y + n³ 6 N 20

40 ric

७५.

H.N	A.N	Date of admit	Date of discharge	Age	Married (Age)	Para	Last (years)	About last (years)	Amenorrhea (month)	Cause of Abortion
148968	52544	2 n. n. 99	5 n. n. 99	22	21	0	0		5	Trauma
148734	52468	2 n. n. 99	4 n. n. 99	32	20	5	$8\frac{1}{2}$	3	1 ⁺	No.
148167	52525	3 n. n. 99	5 n. n. 99	30	19	5	3	0	1 ⁺	No
46742	52516	3 n. n. 99	5 n. n. 99	26	20	2	1	2	3	Infection
148961	52529	3 n. n. 99	5 n. n. 99	29	18	4	1	0	3	No.
148963	52530	3 n. n. 99	5 n. n. 99	38	23	5	1 ⁺	1	5 ⁺	No
148862	52514	3 n. n. 99	6 n. n. 99	25	18	3	$10\frac{1}{2}$	0	3	Trauma
148980	52523	4 n. n. 99	8 n. n. 99	20	18	1	1 ⁺	0	1	Trauma
149116	52610	5 n. n. 99	8 n. n. 99	21					3 ⁺	No
148812	52649	6 n. n. 99	14 n. n. 99	25	24	0	0		6	Trauma
149125	52630	6 n. n. 99	9 n. n. 99	43	20	10	4	2	6	Trauma
149128	52644	6 n. n. 99	8 n. n. 99	19	19	0	0		2	No
149121	52622	8 n. n. 99	11 n. n. 99	22	19	2	1 ⁺	0	6	Trauma
149227	52698	8 n. n. 99	10 n. n. 99	28	18	3	3	2	2	Trauma
149336	52735	8 n. n. 99	14 n. n. 99	27	22	1	5	0	5	Trauma
149243	52728	8 n. n. 99	10 n. n. 99	45	20	11	5	2	2	Trauma
135856	52841	8 n. n. 99	16 n. n. 99	34	18	5	5	1	1	No
128983	52846	8 n. n. 99	12 n. n. 99	21	16	2	2	0	2	No
149241	52826	8 n. n. 99	11 n. n. 99	30	26	3	1 ⁺	1	3 ⁺	Trauma
149219	52667	9 n. n. 99	12 n. n. 99	33	22	4	5	0	3 ⁺	Trauma
148128	52858	9 n. n. 99	11 n. n. 99	24	20	2	$\frac{1}{2}$	0	4	No.
149495	52880	9 n. n. 99	11 n. n. 99	20	19	0	0		5	No
149428	52881	9 n. n. 99	11 n. n. 99	35	22	6	2	2	10	No
149552	52838	10 n. n. 99	15 n. n. 99	35	22	5	5	0	1 ⁺	Trauma
149528	52821	10 n. n. 99	17 n. n. 99	23	23	0	0		3	Trauma
88516	52824	10 n. n. 99	12 n. n. 99	40	18	11	2	0	2	Trauma
149572	52827	10 n. n. 99	12 n. n. 99	37	26	3	3	2	2	Trauma
35677	52808	10 n. n. 99	12 n. n. 99	21	18	4	$\frac{1}{2}$	0		No.
149677	52881	11 n. n. 99	20 n. n. 99	24	18	4	4	1	5	No
149989	52886	11 n. n. 99	15 n. n. 99	35	16	12	1	13	3	Trauma

H.N.	A.N.	Date of Admit	Date of Discharge	Age	Married (Age)	Para	Last (years)	Abort Last (years)	Amniocentesis (month)	Cause of Abortion	
128650	52871	11 M. A. 99	13 M. A. 99	21	18	1	3	0	3	No.	
149627	52875	11 M. A. 99	15 M. A. 99	25	22	3	1	0	2 ⁺	No.	
149691	52882	12 M. A. 99	15 M. A. 99	35	12	6	1 ⁺	0	2 ⁺	Trauma	
149815	53000	14 M. A. 99	16 M. A. 99	28	17	2	8	0	1 ⁺	Trauma	
148490	53010	15 M. A. 99	18 M. A. 99	24	19	3	1	0	2 ⁺	No.	
149819	53014	15 M. A. 99	17 M. A. 99	30	21	2	7	1	4	Trauma	
103266	53033	15 M. A. 99	18 M. A. 99	23	20	1	11 ¹ / ₂	0	1 ¹ / ₂	Trauma	
149701	53039	15 M. A. 99	17 M. A. 99	41	18	8	4	0	4 ⁺	No.	
129063	53041	15 M. A. 99	17 M. A. 99	29	22	1	5 ¹ / ₁₂	0	3	No.	
149932	53044	15 M. A. 99	18 M. A. 99	28	20	3	4	0	4 ⁺	Trauma	
149764	53040	15 M. A. 99	16 M. A. 99	19	15	1	2	1	6 ¹ / ₁₂	Trauma	
149934	53050	16 M. A. 99	19 M. A. 99	26	22	3	1 ⁺	0	4 ⁺	No.	
149999	53085	16 M. A. 99	18 M. A. 99	31	25	7	11 ¹ / ₁₂	0	3 ⁺	No.	
148538	53099	16 M. A. 99	28 M. A. 99	30	26	3	1 ⁺	0	4 ⁺	Infection	
150020	53093	16 M. A. 99	19 M. A. 99	35	21	4	4	1	7	No.	
150152	53153	18 M. A. 99	20 M. A. 99	35	20	7	2 ⁺	4	4 ¹ / ₁₂	2 ⁺	No.
150149	53145	17 M. A. 99	19 M. A. 99	22	18	0		1	3 ¹ / ₁₂	2	No.
84944	53150	17 M. A. 99	20 M. A. 99	28	19	4	1 ⁺	0	3 ⁺	No.	
150153	53158	17 M. A. 99	21 M. A. 99	17	12	0		0	3	Trauma	
150089	53162	17 M. A. 99	21 M. A. 99	23	17	3	11 ¹ / ₁₂	0	5	Trauma	
150056	53118	17 M. A. 99	18 M. A. 99	21	19	0		1	6 ¹ / ₁₂	3 ⁺	Infection Trauma
150148	53143	17 M. A. 99	22 M. A. 99	26	20	1	8	1	1	5	Trauma
149890	53142	18 M. A. 99	20 M. A. 99	20	17	2	11 ¹ / ₁₂	0		3 ⁺	Trauma
149313	53025	18 M. A. 99	20 M. A. 99	38	20	6	5	0		1 ⁺	No.
150220	53183	18 M. A. 99	20 M. A. 99	25	20	1	2	0		3	Trauma
150235	53219	19 M. A. 99	21 M. A. 99	27	18	1	4	0		3	Trauma
150249	53276	19 M. A. 99	22 M. A. 99	22	17	4	1 ⁺	0		4	No.
150241	53207	19 M. A. 99	21 M. A. 99	26	18	4	1 ⁺	0		2 ⁺	Trauma
150241	53221	19 M. A. 99	23 M. A. 99	24	22		8			6	No.
58591	53265	20 M. A. 99	22 M. A. 99	26	20	2	7	0		3	Trauma

H.N	A.N.	Date of Admit	Date of Discharge	Age	Married (Age)	Para	Last (year)	Abort	Last (year)	Amenorrhoea (month)	Cause of Abortion
150538	53280	20.4.99	23.4.99	35	18	8	3	0		6	Trauma
146837	53281	20.4.99	24.4.99	27	21	2	1 ⁺	0		1 ⁺	Chemical (Doubtful)
146837	53333	22.4.99	24.4.99	25	20	2	3	0		3 ⁺	Trauma
85842	53334	22.4.99	24.4.99	33	20	7	1	3	3	2 $\frac{1}{2}$	No
21992	53335	22.4.99	25.4.99	26	10	3	2 ⁺	5	2 $\frac{1}{2}$	1	Infection
150447	53387	23.4.99	25.4.99	40	20	9	2 ⁺	0		1 $\frac{1}{2}$	Trauma
150463	53425	24.4.99	26.4.99	36	24	4	2 $\frac{1}{2}$	0		2	Trauma
150626	53512	26.4.99	28.4.99	27	25	0		1	1 ⁺	1	No
114242	53537	26.4.99	28.4.99	39	27	3	1 ⁺	1	7	?	No
149533	53049	26.4.99	29.4.99	30	30	0		0		6 ⁺	Toxaemia of pregnancy
150566	53517	26.4.99	29.4.99	45	16	7	8	0		2	No
150729	53571	27.4.99	30.4.99	24	18	2	0	2	1 ⁺	2	Trauma
150740	53612	28.4.99	30.4.99	21	21	0		0		2	Trauma
150840	53649	27.4.99	1.5.99	29	19	2	3	0		2 $\frac{1}{2}$	Trauma
150920	53696	30.4.99	1.5.99	43	22	5	12	1	2	2 ⁺	No
150938	53723	31.4.99	1.5.99	30	20	5	14 $\frac{1}{2}$	2	1 ⁺	2	No
147627	51927	19.4.99	11.4.99	28	19	2	1 ⁺	0		4	No
140279	50937	26.4.99	16.4.99	23	16	1	1 ⁺	0		3 ⁺	No
146372	52313	28.4.99	2.4.99	29	22	3	4	0		2	No
148437	52267	27.4.99	4.4.99	29	24	0		2	1 ⁺	3 ⁺	No
86695	52283	27.4.99	3.4.99	25	20	0		0		?	No
148609	52661	28.4.99	9.4.99	25	20	2	2 ⁺	0		3	No
98411	52262	29.4.99	2.4.99	21	18	1	2 ⁺	0		4 ⁺	No
146514	53283	20.4.99	23.4.99	36	18	6	1 ⁺	0	0	6 ⁺	Twins. Polyhydramnios
148765	52488	2.4.99	11.4.99	35		0		4	5	1 $\frac{1}{2}$	No
148864	52504	2.4.99	9.4.99	32	23	3	23 $\frac{1}{2}$	2	2	3	No
149479	52786	9.4.99	11.4.99	19	15	1	1 ⁺	2	2 $\frac{1}{2}$	1 ⁺	Trauma
147857	52928	12.4.99	16.4.99	26	24	0		0		3	Trauma
150116	53104	13.4.99	21.4.99	32	23		2			2	No
148137	52074	26.4.99	26.4.99	34	20	8	1	0		6	No

H.N.	A.N.	Date of Admit	Date of Discharge	Age.	Married (Age)	Para	Last (year)	Abort	Last (year)	Amenorrhea	Cause of Abortion
148132	52092	22 N.V.99	25 N.V.99	22	20	1	1 ⁺	0		1	No
148140	52093	23 N.V.99	26 N.V.99	18	17	0		0		3	No
148259	52146	24 N.V.99	26 N.V.99	32	24	1	1 ⁺	0		1 $\frac{1}{2}$	No
88673	52187	25 N.V.99	28 N.V.99	31	18	2	10 ⁺	3	1 ⁺	1 $\frac{1}{2}$	No
148414	52209	26 N.V.99	28 N.V.99	22	21	0		0		3	No
148418	52221	26 N.V.99	29 N.V.99	22	22	1	$\frac{6}{12}$	0		3	No
148554	52287	27 N.V.99	29 N.V.99	27	22	1	5	1	3	2 ⁺	No
148082	51659	3 N.V.99	19 N.V.99	18	17	0		P		4 ⁺	Trauma
146261	51275	3 N.V.99	7 N.V.99	22	17	1	2	0	0	2 ⁺	No
146374	51315	4 N.V.99	7 N.V.99	30	18	5	2 $\frac{1}{2}$	0		2 $\frac{1}{2}$	No
148562	52574	21 N.V.99	25 N.V.99	22	21	0		0		2	No
148577	52521	23 N.V.99	26 N.V.99	22	20	0		0	1 ⁺	2	Trauma
148580	52614	23 N.V.99	26 N.V.99	22	21	0		0		2	Trauma
148584	52641	27 N.V.99	29 N.V.99	28	26	2	2	0		2 $\frac{1}{2}$	Trauma
148588	52624	30 N.V.99	31 N.V.99	23	22	2	10	2	2	2 ⁺	No
148593	52523	31 N.V.99	31 N.V.99	26	26	0	10	2	2 ⁺	2	No
148597	52523	19 N.V.99	19 N.V.99	21	19	2	1 ⁺	2		4	No
148601	52523	25 N.V.99	26 N.V.99	22	20	1	1 ⁺	0		3 ⁺	No
148622	52462	25 N.V.99	26 N.V.99	23	22	2	4	0		2	No
148623	52462	25 N.V.99	26 N.V.99	23	22	0		0	1 ⁺	0 ⁺	No
148625	52523	26 N.V.99	26 N.V.99	22	20	0		0		1	No
148627	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148628	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148629	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148630	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148631	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148632	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148633	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148634	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148635	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148636	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148637	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148638	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148639	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148640	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148641	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148642	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148643	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148644	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148645	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148646	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148647	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148648	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148649	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148650	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148651	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148652	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148653	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148654	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148655	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148656	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148657	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148658	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148659	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148660	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148661	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148662	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148663	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148664	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148665	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148666	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148667	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148668	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148669	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148670	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148671	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148672	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148673	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148674	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148675	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148676	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148677	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148678	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148679	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148680	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148681	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148682	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148683	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148684	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148685	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148686	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148687	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148688	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148689	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148690	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148691	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148692	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148693	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148694	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148695	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148696	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148697	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148698	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No
148699	52523	26 N.V.99	26 N.V.99	22	21	0		0		2	No

Case abortion 2499 5 months 82 no.
 18 months 18 no.
 100 no.

Inflow 100 no. Inevitable abortion 90 no. - Incomplete abort: 82
 Complete abort 8
 Threatened abortion 10 no.

∴ Inevitable abortion 90% Incomplete abortion 82%
 Complete abortion 8%
 Threatened abortion 10%

Cause of Abortion. 100 no.

Trauma 41 %
 Infection 3 %
 Unknown cause 53 %
 Toxaemia pregnancy 1 % - edema, high B.P., abnormal
 Twins & polyhydramion 1 %
 Chemical (drugs) 1 % Criminal abortion

Threatened abortion 10 no. - Unknown cause 4 no. = 40%
 Infection 2 no. = 20%
 Trauma 1 no. = 10%
 Some of Inf. hepatitis

Inevitable abortion 90 no. Unknown cause 46 no. = 51.1%
 Trauma 40 no. = 44.4%
 Some of Infection 1 no. = 1.1%
 Toxaemia preg. 1 no. = 1.1%
 Twins & polyhydramion 1 no. = 1.1%
 Chemical 1 no. = 1.1%

John Jones
 1914

B.P.	Heart	Lung	Bl.K.	Urine		Hb	Rbc	W.b.c	Polyp	lym	Type of Abortion	Rx	Result
				Alb	Sugar								
140/70	O.K.	O.K.	neg	neg	neg	40	240	6200	86	14	Complete	Symptomatic	Improved
110/80	O.K.	O.K.	neg	neg	neg	44	375	5050	88	23	Incomplete (Septic)	Curette Antibiotic	Improved
108/60	-	-	neg	neg	neg	44	395	6450	82	18	Incomplete	Curette	Improved
110/70	-	-	neg	neg	2+ (first 2 days)	64	265	8400	62	38	Incomplete	Curette	Improved
120/80	O.K.	O.K.	-	-	-	44	365	8250	80	16	Incomplete	Curette	Improved
100/60	-	-	-	-	-	44	375	10600	86	24	Incomplete	Curette	Improved
90/60	O.K.	O.K.	-	-	-	54	325	6250	83	28	Incomplete (Septic)	Curette Antibiotic	Improved
110/70	-	-	-	neg	neg	68	459	13400	68	32	Incomplete	Curette	Improved
120/80	O.K.	O.K.	neg	-	-	-	-	-	-	-	Incomplete	Curette	Improved
180/142	O.K.	O.K.	neg	1+	neg	48	386	8600	82	28	Complete	Analgesic	Improved
130/90	-	-	neg	neg	neg	64	383	5850	85	25	Incomplete	Curette	Improved
110/80	O.K.	O.K.	neg	neg	neg	44	335	5100	94	6	Incomplete (Habitual)	Curette	Improved
110/70	O.K.	O.K.	neg	neg	neg	68	295	12000	71	29	Incomplete	Curette	Improved
120/86	Extrem systolic	O.K.	neg	neg	neg	54	241	8850	93	8	Incomplete	Curette	Improved
130/90	O.K.	O.K.	-	-	-	61	341	6200	62	20 E	Inevitable abor	Curette	Improved
110/80	O.K.	O.K.	-	-	-	58	315	6600	86	14	Incomplete (Septic)	Curette Antibiotic	Improved
150/110	-	-	-	-	-	68	389	10600	83	12 E	Incomplete	Curette	Improved
130/80	O.K.	O.K.	neg	neg	neg	51	140	4800	85	14 1	Threatened	Sedation Bl. Trans.	Improved
110/80	-	-	neg	neg	neg	51	135	6800	93	8	Incomplete	Curette	Improved
110/70	-	-	neg	neg	neg	78	415	2650	51	31 E	Incomplete	Curette	Improved
128/80	-	-	neg	neg	neg	78	390	4400	38	69	Threatened	Symptom	Improved
110/80	-	-	neg	neg	neg	44	296	7400	78	22 E	Threatened	Symptom	Improved
110/70	-	-	neg	neg	neg	48	315	8800	64	36	Incomplete	Curette	Improved
100/60	Rheumatic Mitral stenosis	O.K.	neg	neg	neg	51	285	5800	76	24	Incomplete	Curette	Improved
140/90	O.K.	O.K.	neg	neg	2+	81	365	11500	84	13	Incomplete (Habitual)	Curette	Improved
114/90	O.K.	O.K.	-	neg	neg	51	-	-	-	-	Incomplete	Curette	Improved
110/70	-	-	-	neg	neg	-	-	-	-	-	Incomplete	Curette	Improved
120/80	O.K.	O.K.	neg	neg	neg	81	441	6000	84	16	Incomplete	Curette	Improved
112/70	O.K.	O.K.	-	-	-	-	-	-	-	-	Incomplete	Curette	Improved
110/70	-	-	neg	-	-	44	396	6900	88	22	Threatened	Sedation	Improved

B.P.	Heart	Lung	Bl. K.	Urine		Hb	Rbc.	Wbc	Polyp	lym.	Type of Abortion	Rx	Result.
				Alb	Sugar								
110/80	O.K.	O.K.		neg	neg	54	354	5350	48	52	Incomplete	Curette	Improved
120/80	-	-	neg	neg	neg	34	2,10	5450	83	18	Incomplete (Septic)	Curette Antibiotics	Improved
100/60	O.K.	O.K.	neg	neg	4 ⁺	48	2,40	4800	20	18 ^E ₁₂	Incomplete	Curette	Improved
90/60	O.K.	O.K.	neg	-	-	39	2,80	5300	80	20	Incomplete	Curette	Improved
110/40	O.K.	O.K.	-	neg	neg		2,55	6250	80	12	Incomplete	Curette	Improved
90/60	O.K.	O.K.	neg	-	-	40	2,30	8350	28	22	Incomplete	Curette	Improved
120/80	O.K.	O.K.	neg	neg	2 ⁺	81	3,38	4000	29	20 ^E ₁	Incomplete	Curette	Improved
110/80	O.K.	O.K.	neg	-	-	40	2,05	6950	29	21	Incomplete	Curette	Improved
108/80	Rheumatic H.D.	O.K.	-	-	-	-	-	-	-	-	Incomplete	Curette	Improved
104/80	O.K.	O.K.	neg	neg	neg	58	2,80	14900	84	16	Incomplete	Curette	Improved
120/80	O.K.	O.K.	neg	-	-	28	3,81	6600	29	21	Incomplete	Curette	Improved
120/100	O.K.	O.K.	neg	neg	neg	54	2,59	6050	27	21 ^E ₂	Threatened	Supportive Symptomatic	Improved
120/80	O.K.	O.K.	sp ⁺	neg	2 ⁺	54	2,48	5650	20	30	Incomplete	Curette	Improved
110/80	O.K.	O.K.	neg	neg	neg	51	2,60	5200	28	23	Threatened	Sedative	Improved
110/40	O.K.	O.K.	neg	neg	neg	38	2,24	11000	83	18	Incomplete (Septic)	Antibiotics Curette	Improved
80/40	O.K.	O.K.	neg	neg	neg	39	2,99	6450	22	28	Incomplete	Curette	Improved
96/60	O.K.	O.K.	neg	-	-	68	3,91	13950	28	22	Incomplete (Septic)	Curette Antibiotics	Improved
120/80	-	-	neg	neg	neg	68	3,40	5600	59	35 ^E ₆	Incomplete	Curette	Improved
100/60	O.K.	O.K.	neg	neg	neg		2,30	10200	82	18	Incomplete	Curette	Improved
130/90	O.K.	O.K.	neg	neg	neg	58	2,59	26050	84	16	Incomplete (Septic)	Curette Antibiotics	Improved
90/60	O.K.	O.K.				51	2,88	12650			Incomplete	Curette	Improved
106/60	O.K.	O.K.	neg			64	3,10	8000	84	16	Incomplete (Septic)	Curette Antibiotics	Improved
120/80	O.K.	O.K.	neg	neg	neg	51	3,05	4850	22	21 ^E ₈	Incomplete	Curette	Improved
120/80	O.K.	O.K.	neg	neg	neg	28	3,21	8220	28	21 ^E ₁	Incomplete	Curette	Improved
110/80	O.K.	O.K.	neg	neg	neg	61	3,15	5550	84	16	Incomplete	Curette	Improved
100/60	-	-	neg			64	2,50	4000	55	45	Incomplete	Curette	Improved
100/60	-	-	neg	neg	neg	74	3,60	9400	82	18	Incomplete	Curette	Improved
98/60	-	-	neg	neg	neg	64	3,96	3900	85	13	Incomplete	Curette	Improved
84/80			neg	neg	neg	64	3,00	13200	86	14	Complete	Symptomatic	Improved
120/60	O.K.	O.K.	neg	neg	neg	28	3,85	5500	29	20 ^E ₁	Incomplete	Curette	Improved

D.P. (mm Hg)	Heart	Lung	Bl. K.	Urine		Hb %	R.b.c. Million	W.b.c.	Polyph	Lym	Type of Abortion	Rx	Result.
				Alb	Sugar								
120/80	O.K.	O.K.									Complete	Symptomatic	Improved
100/60	O.K.	O.K.				61	2,50	5250	75	25	Incomplete	Curette	Improved
130/90	O.K.	O.K.				39	2,12	7000	66	34	Threatened	Sedative Hormone	Improved
100/60	O.K.	O.K.		neg	neg	68	3,05	6200	74	26	Threatened	Sedative	Improved
130/90	O.K.	O.K.	neg	neg	neg						Incomplete (Septic)	D+C Anti biotics	Improved
110/80	O.K.	O.K.	-	neg	neg	61	2,40	7200	72	28	Incomplete	Curette	Improved
100/60	O.K.	O.K.	neg	neg	neg	57	3,05	6000	58	42	Incomplete	Curette	Improved
140/100	O.K.	O.K.	neg	neg	1+	74	3,72	5800	86	9 (E5)	Incomplete	Curette	Improved
120/80	O.K.	O.K.	neg	neg	neg	68	2,96	4900	60	35 ^E / ₅	Incomplete	Curette	Improved
110/80	O.K.	O.K.	neg	neg	neg	74	2,91	10200	85	15	Complete	Symptomatic	Improved
94/76	O.K.	O.K.	neg	neg	neg	39	2,05	6600	86	14	Complete	Oxytocic	Improved
120/90	O.K.	O.K.	-	neg	neg	68	2,60	7200	68	18 ^E / ₅	Incomplete	Curette	Improved
120/90	O.K.	O.K.	-	neg	neg	58	2,40	4000	72	28	Threatened	Sedative	Improved
110/80	O.K.	O.K.	2+	neg	neg						Incomplete (Septic)	Curette Anti biotics	Improved
120/80	O.K.	O.K.	neg			68	3,64	10630	85	15	Complete	Oxytocic	Improved
100/80	O.K.	O.K.	neg	neg	neg	74	3,65	7350	58	43	Incomplete	Curette	Improved
80/62	O.K.	O.K.	neg	neg	neg	58	2,30	7800	69	31	Incomplete	Curette	Improved
140/100	O.K.	O.K.	neg			78	3,00	8400	69	28 ^E / ₄	Incomplete	Curette	Improved
104/80	-	-	neg	neg	1+						Incomplete	Curette	Improved
116/60	-	-	neg	neg	neg	54	3,00	13600	95	5	Incomplete	Curette	Improved
130/90	O.K.	O.K.	4+	neg	neg	68	2,39	8900	74	24 ^E / ₂	Incomplete	Curette	Improved
110/80	-	-	neg	neg	neg	54	3,10	8100	82	18 ^E / ₁	Incomplete	Curette	Improved
160/90	O.K.	O.K.	neg	neg	neg	48	3,75	8000	62	38	Incomplete	Curette	Improved
110/80	O.K.	O.K.	neg	-	-	60	2,15	7600	88	12	Incomplete (Septic)	Curette Anti biotics	Improved
90/60	O.K.	O.K.	neg	-	-	68	3,10	5600	87	12 ^E / ₁	Incomplete	Curette	Improved
130/90	O.K.	O.K.	-	neg	neg	74	3,20	6800	59	26 ^E / _{1,14}	Incomplete	Curette	Improved
100/80	O.K.	O.K.	-	-	-						Incomplete	Curette	Improved
100/80	O.K.	O.K.	neg	neg	neg	68	3,15	5000	76	21 ^E / ₃	Incomplete	Curette	Improved
100/80	O.K.	O.K.	neg	neg	neg	39	2,00	6800	67	33	Threatened	Sedative	Improved
110/80	O.K.	O.K.	4+	neg	neg	78	3,10	12400	78	22	Incomplete	Curette	Improved

เอกสารส่วนบุคคล ศาสตราจารย์นายแพทย์เสม พริ้งพวงแก้ว

สพ 1

2/1.4 สถิติสถิติรวมการแต่งศพ ช่วงพ.ศ. 2494 (ชื่อเรื่อง)
- 2499 โรงพยาบาลเดนิม

พ.ศ. 2494 - 2499

จำนวน 2 แผ่น

แฟ้มที่ 2

กล่องที่ 1

Bleeding in Pregnancy exclusive of abortion: Incidence 1:80

สถิติสักรวม พ.ศ. ๒๔๘๔-๒๔๘๕

จำนวนคนไข้มาก (พ.ศ. ๒๔๘๔-๒๔๘๕)	๑๘๕๕๑	คน
คนไข้มีอาการเลือดออกก่อนคลอด(ระยะครรภ์ตั้งแต่ ๖ เดือนขึ้นไป)	๒๓๓	"
คิดเป็นอัตราส่วน	๑ : ๘๐ (๘๑.๕)	
คนไข้มีอาการเลือดออกมากหลังคลอด	๓๐๑	"
คิดเป็นอัตราส่วน	๑ : ๖๐ (๖๒.๔)	

= 1.2%
= 1.6%

.....

คนไข้มีอาการเลือดออก	๒๓๓	คน
Placenta previa (Baylor University Hospital 32305 Deliveries - Placenta previa 137 cases incidence = 0.42%)	๑๘๕	ราย = 81.78%
Abruptio placenta	๔๘	" = 19.22%
คิดอัตราส่วน Placenta previa	๑ : ๑๐๐	
" Abruptio placenta	๑ : ๔๓๐	

อัตราตาย

แม่	(Placenta previa)	๓.๑ %
เด็ก	(" ")	๒.๔ %
แม่	(Abruptio placenta)	๒.๒ %
เด็ก	(" ")	๑.๐๐ %
แม่	(Post partum hemorrhage)	๒.๓ %

.....

Incidence

① Bleeding in Pregnancy (not including abortion) = 1:80 = 1.2%

② Abortion 932:

8619: 932 = 932: 8619
Total Deliveries: abortion 1:9 = 10%
abortion: Normal Deliveries

③ Post-partum bleeding 1:60 = 1.6%

ABORTION

๑ มกราคม ๒๔๘๔ ถึง ๓๑ ธันวาคม ๒๔๘๔

No.	Abortions.	จำนวน	คิดเป็นเปอร์เซ็นต์ %
1.	Incomplete abortion	๒๔๔	๒๔.๐๔ %
2.	Complete abortion	๑๒๓	๑๒.๑๔ %
3.	Threatened abortion	๔๐	๔.๕๔ %
4.	Inevitable abortion	๗๖	๘.๑๕ %
5.	Criminal abortion	๖	๐.๖๔ %
6.	Habitual abortion	๑	๐.๑ %
7.	Therapeutic abortion	๑	๐.๑๐ %
8.	Missed abortion	๑	๐.๑๐ %
	Total cases	๔๓๖	—

CAUSES OF EVERY KINDS OF ABORTIONS.

1. INCOMPLETE ABORTION.

No.	สาเหตุ	จำนวน	คิดเป็นเปอร์เซ็นต์ %
1.	ไม่ทราบสาเหตุ	๔๒๒	๔๑.๐๕ %
2.	หกล้ม	๒๖	๑๐.๒๔ %
3.	รูดกระแทก	๑๓	๒.๐๑ %
4.	บกพร่องพันธุ	๑๕	๒.๓๓ %
5.	ตกมันโค	๑๗	๒.๖๓ %
6.	ตกเก้าอี้	๑	๐.๑๐ %
7.	กินยา	๔	๐.๖๒ %
8.	ร่วมประเวณี	๑	๐.๑๐ %
9.	ไร่	๕	๐.๗๑ %

2. COMPLETE ABORTION.

No.	สาเหตุ	จำนวน	คิดเป็นเปอร์เซ็นต์ %
1.	ไม่ทราบสาเหตุ	๑๐๓	๔๓.๗๓ %
2.	หกล้ม	๑๓	๑๐.๕๖ %
3.	รูดกระแทก	๒	๑.๖๒ %
4.	บกพร่องพันธุ	๓	๒.๔๓ %
5.	ตกมันโค	๒	๑.๖๒ %

3. THREATENED ABORTION.

No.	สาเหตุ	จำนวน	คิดเป็นเปอร์เซ็นต์
1.	ไม่ทราบสาเหตุ	๓๑	๘๘.๓๕ %
2.	รูดกระแทก	๒	๒.๕ %
3.	ยกของหนัก	๒	๒.๕ %
4.	ตกบันได	๑	๑.๒๕ %
5.	หกล้ม	๔	๕.๐ %

4. INEVITABLE ABORTION.

No.	สาเหตุ	จำนวน	คิดเป็นเปอร์เซ็นต์
1.	ไม่ทราบสาเหตุ	๖๔	๘๔.๒๑ %
2.	หกล้ม	๕	๖.๕๓ %
3.	รูดกระแทก	๒	๒.๖๓ %
4.	ยกของหนัก	๑	๑.๓๑ %
5.	ตกบันได	๓	๓.๘๔ %
6.	ตกเก้าอี้	๑	๑.๓๑ %

5. CRIMINAL ABORTION.

No.	สาเหตุ	จำนวน	คิดเป็นเปอร์เซ็นต์
1.	ไม่ทราบสาเหตุ	๓	๕๐.๐๐ %
2.	Schizophrenia.	๑	๑๖.๖๖ %
3.	Drugs.	๒	๓๓.๓๓ %

6. HABITUAL ABORTION.

No.	สาเหตุ	จำนวน	คิดเป็นเปอร์เซ็นต์
1.	ไม่ทราบสาเหตุ	๑	๑๐๐ %

7. THERAPEUTIC ABORTION.

No.	สาเหตุ	จำนวน	คิดเป็นเปอร์เซ็นต์
1.	T.B.	๑	๑๐๐ %

8. MISSED ABORTION.

No.	สาเหตุ	จำนวน	คิดเป็นเปอร์เซ็นต์
1.	ไม่ทราบสาเหตุ	๑	๑๐๐ %

เอกสารส่วนบุคคล ศาสตราจารย์นายแพทย์เสมอ พริ้งพวงแก้ว

สพ 1

2/1.5

ศึกษาโดยโรงพยาบาลในภาคเหนือ (ชื่อเรื่อง)

ช่วง พ.ศ. 2494 - 2500 โรงพยาบาล

พ.ศ. 2494-2500

จำนวน 23 แผ่น

แฟ้มที่ 2

กล่องที่ 1





Site 4

Following

Pytho anthon
7/13/94
D. anthon
9.0.95
D. anthon
D. anthon

5/22/95

5/28/95

28.0.95

DRD

13.0.0

6.0.0.0

Pytho anthon

Pytho anthon

10.5.0.0

10.8.0.0

19.0.96

DRD

2494

2499

2498.

24952

2496

2499

2497.

2499

S

2499

2499

2499

2499

2499

2499

2499

2499

2499

2499

2499

2499

2499

2499

2499

2499

2499

2499

2499

2499

2499

2499

2499

2499

2499

2499

2498

2498

2498

2498

2498

2498

2498

2498

2498

2498

2498

2498

2498

2498

2499 (10)

XVI

XVI

er hoo G.
Wm Peppic mles

XVII

XVII

Repealed
1800



407

407

41

31

416

41

41

41

XV

XVI

XVII

XVIII

1499

XIX

XX

XXI

Hydrom +
lily pond of
dandelion

Carbide
200 g at 10/12

Hydrom +
oil 6. canol

No. F. d.

about
100 g. and
4-5 g

acute 10/12

500 g. 2500
and 1/2 cup
m. 4. 0/12 300 g.

Hydrom
(X-ray)

(X-ray)

100 g. 2500

DEAD
2500-2000

(1.6)

DEAD
2000-2500

Hydrom

6/6 cur. -
Cardiac
pulser
(X-ray)

NOFU

DEAD
Q.W.A. 2500

months P.O.

Cardia

Pylosic
(Yrmy)

Pyburn

Mr F. U.

locus 25

1 Dec 20)
metastatic
Ca. to mesenteric
lymph glands.

Pyloric
part.

ตาราง B จำนวนผู้สูงอายุที่มีอายุ 65 ปีขึ้นไป

อายุ (ปี)	จำนวนผู้สูงอายุ	จำนวนผู้สูงอายุ
11 - 20	—	4
21 - 30	8	9
31 - 40	2	7
41 - 50	4 (3)	15 (24)
51 - 60	11 *	11
61 - 70	7	2
71 - 80	—	2

รวมผู้สูงอายุที่มีอายุ 65 ปีขึ้นไป 215
 รวมผู้สูงอายุ 305
 * รวมผู้สูงอายุที่มีอายุ 51-60 ปี

จำนวน	จำนวน	จำนวน
จำนวน	30 (31)	86
จำนวน	4	11
จำนวน	1	

ตาราง E จำนวนผู้สูงอายุที่มีอายุ 65 ปีขึ้นไป

รวมผู้สูงอายุ	รวม	
รวมผู้สูงอายุ	205	
รวมผู้สูงอายุ	รวม	รวมผู้สูงอายุ 365
รวมผู้สูงอายุ	5	
1-6 ปี	21	58.88
7-12 ปี	7	
2-6 ปี	1	
7-12 ปี	—	
13-19 ปี	— (1)	
20-36 ปี	1	

กระดาดสร้าง

ผู้ร่าง

วันที่

ผู้ตรวจ

วันที่

เรื่อง

ตรวจ ๕ ลักษณะที่พบในกระดาด

ตำแหน่ง	จำนวน	หมายเหตุ
Cardia	3	
Greater Curvature	5	(5) ①
Lesser Curvature	5	
Pylorus	10	(5)
Pylorus + Duodenum	4	
ส่วนหัว ๑ หรือ 2/3	3	(2)
ท้อง IV หรือ กระเพาะอาหารส่วนล่าง region ๔, ๕	1	
โดยทั่วทั้งกระเพาะ	9	
หมายเหตุ: pylorus ตรวจพบ ลักษณะโดยทั่วทั้งกระเพาะ		

ตรวจ ๕ ลักษณะที่พบในกระดาด

ตำแหน่ง	จำนวน	หมายเหตุ
ส่วนหัว	7 (8)	19.95 12.14.1
ส่วนกลาง	5	
ส่วนล่าง	3	
ส่วนหัว + ส่วนกลาง	12	37.05 31.14.1
กระเพาะ + ส่วนหัว (H. l. m.)	1	
กระเพาะ + ส่วนกลาง (H. m. m.)	1	
กระเพาะ + ส่วนล่าง	2	
กระเพาะ + ส่วนหัว + ส่วนกลาง	1	
กระเพาะ + ส่วนหัว + ส่วนล่าง	2	

Summary of Gastric Analysis

Clava Gastric Analysis

Hypo-acidity

จำนวน

14 13

Hyper-acidity

4

Total

18

Summary

จำนวน

จำนวน

จำนวน

Gastro-jejunostomy

3

Gastrostomy

2

Subtotal Gastrostomy + Gastro-jejunostomy

9

Partial esophago-gastrostomy & anastomosis

1

Exploratory + Biopsy

5

Total

16 15

Summary of Gastric Surgery

จำนวน

จำนวน

Chronic inflammatory reaction - no evidence of

3

Malignant cells in gastric fluid

1

* Adeno-carcinoma

7

Infiltrating adenocarcinoma

1

Signet cell carcinoma

2

Mucoid carcinoma

1

Hypertrophic Gastritis

1

Metastatic Adenocarcinoma

1

Total

19 18

* 1 case of adenocarcinoma on top of peptic ulcer.

① under word adenocarcinoma . 207

กระต่ายร่าง

ผู้ร่าง

วันที่

ผู้ตรวจ

วันที่

เรื่อง

มคอ ๑๐ Metastasis

ลักษณะ metastasis	จำนวน
Duodenum + Regional lymph gland	1
Regional lymph gland + Omentum	8
Liver	3
Ovary	1
Testis/ovary	3

ของมะเร็ง

ค่า	เฉลี่ย	ค่าจริง
ค่าเฉลี่ยของค่าเฉลี่ย	9	25.7 25
ค่าเฉลี่ยของค่าเฉลี่ย (ค่าเฉลี่ยของค่าเฉลี่ย)	10.9	29.8
ค่าเฉลี่ย	15	42.8 41.7
ค่าเฉลี่ย	2	5.6

มคอ ๑๐ Survival rate in 5 year

analyse both metastasis in O.P.D and to find

16 re Discharge metastasis to find to find to find

1 re metastasis to metastasis 1, 9 mo PO

1 re metastasis

มคอ ๑๐ Discharge metastasis

one metastasis to metastasis	2 re
one metastasis to metastasis	3 re
one metastasis to metastasis	2 re
Testis/ovary metastasis	8 re

1 re metastasis to metastasis 1, 9 mo PO
1 re metastasis to metastasis 1, 9 mo PO
1 re metastasis to metastasis 1, 9 mo PO

[illegible]

၁၂:၁၆၀ နှစ် ၁၁၀၀ ဟု ၁၂:၁၆၀ နှစ် ၁၁၀၀ ၃၆:၅၉ ၁၆၀ ၁:၁.၆၄

[illegible]

Handwritten notes at the bottom of the page:

10/6/1957
10/6/1957

20. 10. 35 21. 10. 35 22. 10. 35 23. 10. 35 24. 10. 35 25. 10. 35 26. 10. 35 27. 10. 35 28. 10. 35 29. 10. 35 30. 10. 35 31. 10. 35 32. 10. 35 33. 10. 35 34. 10. 35 35. 10. 35 36. 10. 35 37. 10. 35 38. 10. 35 39. 10. 35 40. 10. 35 41. 10. 35 42. 10. 35 43. 10. 35 44. 10. 35 45. 10. 35 46. 10. 35 47. 10. 35 48. 10. 35 49. 10. 35 50. 10. 35 51. 10. 35 52. 10. 35 53. 10. 35 54. 10. 35 55. 10. 35 56. 10. 35 57. 10. 35 58. 10. 35 59. 10. 35 60. 10. 35 61. 10. 35 62. 10. 35 63. 10. 35 64. 10. 35 65. 10. 35 66. 10. 35 67. 10. 35 68. 10. 35 69. 10. 35 70. 10. 35 71. 10. 35 72. 10. 35 73. 10. 35 74. 10. 35 75. 10. 35 76. 10. 35 77. 10. 35 78. 10. 35 79. 10. 35 80. 10. 35 81. 10. 35 82. 10. 35 83. 10. 35 84. 10. 35 85. 10. 35 86. 10. 35 87. 10. 35 88. 10. 35 89. 10. 35 90. 10. 35 91. 10. 35 92. 10. 35 93. 10. 35 94. 10. 35 95. 10. 35 96. 10. 35 97. 10. 35 98. 10. 35 99. 10. 35 100. 10. 35

[Faint handwritten notes at the bottom of the page, possibly from another document or bleed-through.]

(322) α $\frac{32.051}{22.81}$ $\frac{11.01}{11.01}$ $\frac{11.01}{11.01}$ (322) $\frac{22.81}{11.01}$

of hypoxia

missing transaction in Subtotal & extracting value

Handwritten text on a piece of aged paper, possibly a label or note, with some ink bleed-through from the reverse side. The text is written in cursive and includes the words "Inoperable" and "Gastro-jejunostomy".

11/10 more info in Wopsy 15th in Palliative
of 11/10 24 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038

до Лейпцига 10 м (~~27.7.1911~~)

Qumranite was the author of the book of Jubilees.

nam wuduun loga kani wudu: 180 515

Описание 35 н.б.: 15 н.б. в музее, 10 н.б. в архиве, 10 н.б. в библиотеке, 10 н.б. в частной коллекции.

regional lymph gland n= omentum (899 n= 22.24%)

1890
 1891
 1892
 1893
 1894
 1895
 1896
 1897
 1898
 1899
 1900
 1901
 1902
 1903
 1904
 1905
 1906
 1907
 1908
 1909
 1910
 1911
 1912
 1913
 1914
 1915
 1916
 1917
 1918
 1919
 1920
 1921
 1922
 1923
 1924
 1925
 1926
 1927
 1928
 1929
 1930
 1931
 1932
 1933
 1934
 1935
 1936
 1937
 1938
 1939
 1940
 1941
 1942
 1943
 1944
 1945
 1946
 1947
 1948
 1949
 1950
 1951
 1952
 1953
 1954
 1955
 1956
 1957
 1958
 1959
 1960
 1961
 1962
 1963
 1964
 1965
 1966
 1967
 1968
 1969
 1970
 1971
 1972
 1973
 1974
 1975
 1976
 1977
 1978
 1979
 1980
 1981
 1982
 1983
 1984
 1985
 1986
 1987
 1988
 1989
 1990
 1991
 1992
 1993
 1994
 1995
 1996
 1997
 1998
 1999
 2000
 2001
 2002
 2003
 2004
 2005
 2006
 2007
 2008
 2009
 2010
 2011
 2012
 2013
 2014
 2015
 2016
 2017
 2018
 2019
 2020
 2021
 2022
 2023
 2024
 2025
 2026
 2027
 2028
 2029
 2030
 2031
 2032
 2033
 2034
 2035
 2036
 2037
 2038
 2039
 2040
 2041
 2042
 2043
 2044
 2045
 2046
 2047
 2048
 2049
 2050
 2051
 2052
 2053
 2054
 2055
 2056
 2057
 2058
 2059
 2060
 2061
 2062
 2063
 2064
 2065
 2066
 2067
 2068
 2069
 2070
 2071
 2072
 2073
 2074
 2075
 2076
 2077
 2078
 2079
 2080
 2081
 2082
 2083
 2084
 2085
 2086
 2087
 2088
 2089
 2090
 2091
 2092
 2093
 2094
 2095
 2096
 2097
 2098
 2099
 2100
 2101
 2102
 2103
 2104
 2105
 2106
 2107
 2108
 2109
 2110
 2111
 2112
 2113
 2114
 2115
 2116
 2117
 2118
 2119
 2120
 2121
 2122
 2123
 2124
 2125
 2126
 2127
 2128
 2129
 2130
 2131
 2132
 2133
 2134
 2135
 2136
 2137
 2138
 2139
 2140
 2141
 2142
 2143
 2144
 2145
 2146
 2147
 2148
 2149
 2150
 2151
 2152
 2153
 2154
 2155
 2156
 2157
 2158
 2159
 2160
 2161
 2162
 2163
 2164
 2165
 2166
 2167
 2168
 2169
 2170
 2171
 2172
 2173
 2174
 2175
 2176
 2177
 2178
 2179
 2180
 2181
 2182
 2183
 2184
 2185
 2186
 2187
 2188
 2189
 2190
 2191
 2192
 2193
 2194
 2195
 2196
 2197
 2198
 2199
 2200
 2201
 2202
 2203
 2204
 2205
 2206
 2207
 2208
 2209
 2210
 2211
 2212
 2213
 2214
 2215
 2216
 2217
 2218
 2219
 2220
 2221
 2222
 2223
 2224
 2225
 2226
 2227
 2228
 2229
 2230
 2231
 2232
 2233
 2234
 2235
 2236
 2237
 2238
 2239
 2240
 2241
 2242
 2243
 2244
 2245
 2246
 2247
 2248
 2249
 2250
 2251
 2252
 2253
 2254
 2255
 2256
 2257
 2258
 2259
 2260
 2261
 2262
 2263
 2264
 2265
 2266
 2267
 2268
 2269
 2270
 2271
 2272
 2273
 2274
 2275
 2276
 2277
 2278
 2279
 2280
 2281
 2282
 2283
 2284
 2285
 2286
 2287
 2288
 2289
 2290
 2291
 2292
 2293
 2294
 2295
 2296
 2297
 2298
 2299
 2300
 2301
 2302
 2303
 2304
 2305
 2306
 2307
 2308
 2309
 2310
 2311
 2312
 2313
 2314
 2315
 2316
 2317
 2318
 2319
 2320
 2321
 2322
 2323
 2324
 2325
 2326
 2327
 2328
 2329
 2330
 2331
 2332
 2333
 2334
 2335
 2336
 2337
 2338
 2339
 2340
 2341
 2342
 2343
 2344

~~10 - 2 m~~

mortality rate 1500000 13 no one 2 no current case

မှ.နှစ် ၁၀ ပုံ ဖြစ်သည်) ဖြစ်ပါသည်။ Mortality rate ဖြစ်သည် = 15.4

low = 285 (in 35 total) us\$ @ 90 mm on Vah von mich

Five years survival rate 100% with cystic disease 1970

Table II. Operations

Survive -	Survive operation (on stomach)	13	4 persons
(20)	operation (on stomach)	4	
17	operation (on stomach)	4	
one 15	operation 10 on stomach	2	
	operation (on stomach)	9	
	operation 10 on stomach	4	
operation		3	

Mortality rate operation - 10 on stomach 10 on stomach

Operation	No. Survive op ²	%	Dead	%
operation	3			
operation	8		1	
	2			
operation & operation	4		1	

Op² 10 on stomach 2 - M. R. 15.4.1.

(Lecture 10) 25-11-19

20/11

การผ่าตัดของ SURGICAL CASES of Stomach. 20 CASES.

วันที่ 6 พฤษภาคม 2445 - 11 มิถุนายน 2445.

1. Peptic Ulcer. 13 - 18

2. Carcinoma of Stomach 7 cases 35%.

การผ่าตัด Gastrectomy & Gastrojejunostomy 18 cases

การผ่าตัด Gastrostomy 1 case

On 1 case of Stomach cancer

Peptic Ulcer

Carc. of Stomach

26 - 57

26 - 70

average

45

48

เมื่อแรกมีอาการเป็นมา 1 ปี 2 เดือน 3 เดือน 4 เดือน 5 เดือน 6 เดือน 7 เดือน 8 เดือน 9 เดือน 10 เดือน 1 ปี 2 ปี 3 ปี 4 ปี 5 ปี 6 ปี 7 ปี 8 ปี 9 ปี 10 ปี

อาการที่พบ

1. อาการปวดท้อง

2. อาการเบื่ออาหาร

3. อาการคลื่นไส้

4. อาการอาเจียน

5. อาการท้องอืด

6. อาการน้ำหนักลด

7. อาการโลหิตในอุจจาระ

8. อาการซีด

9. อาการอ่อนเพลีย

10. อาการท้องบวม

11. อาการท้องผูก

12. อาการท้องเสีย

13. อาการท้องจุก

14. อาการท้องเฟ้อ

15. อาการท้องอืด

16. อาการท้องบวม

17. อาการท้องผูก

18. อาการท้องเสีย

19. อาการท้องจุก

20. อาการท้องเฟ้อ

[illegible]

CASE	NAME	AGE	SEX	YEAR RIDGE	OPERATIONAL	PAST HISTORY	PHYSICAL EXAM.	LAB FINDING	PAST HISTORY	PROVISIONAL DIAG.	X-RAY	GASTRIC ANALYSIS	PE OPERATION	MICROSCOPE EXAM.	DIAG.
1000	JOHN D. BROWN	47	Q	7	1914	no previous epigastric pain	no definite finding	Hb 54% Rbc 2.4M Wbc 5500 Eos 2% Sed 1000	no	Peptic ulcer	no	Hyperacidity	Gastrectomy	definitely Ulcer (Long ulcers free ulcer)	249
1001	JOHN D. BROWN	33	Q	8	1915	no previous epigastric pain	no definite finding	Hb 40% Rbc 2.4M Wbc 5500 Eos 2% Sed 1000	no	Peptic ulcer	no	Hyperacidity	Gastrectomy	definitely Ulcer (Long ulcers free ulcer)	249
1002	JOHN D. BROWN	47	Q	7	1914	no previous epigastric pain	no definite finding	Hb 54% Rbc 2.4M Wbc 5500 Eos 2% Sed 1000	no	Peptic ulcer	no	Hyperacidity	Gastrectomy	definitely Ulcer (Long ulcers free ulcer)	249
1003	JOHN D. BROWN	47	Q	7	1914	no previous epigastric pain	no definite finding	Hb 54% Rbc 2.4M Wbc 5500 Eos 2% Sed 1000	no	Peptic ulcer	no	Hyperacidity	Gastrectomy	definitely Ulcer (Long ulcers free ulcer)	249
1004	JOHN D. BROWN	47	Q	7	1914	no previous epigastric pain	no definite finding	Hb 54% Rbc 2.4M Wbc 5500 Eos 2% Sed 1000	no	Peptic ulcer	no	Hyperacidity	Gastrectomy	definitely Ulcer (Long ulcers free ulcer)	249
1005	JOHN D. BROWN	47	Q	7	1914	no previous epigastric pain	no definite finding	Hb 54% Rbc 2.4M Wbc 5500 Eos 2% Sed 1000	no	Peptic ulcer	no	Hyperacidity	Gastrectomy	definitely Ulcer (Long ulcers free ulcer)	249
1006	JOHN D. BROWN	47	Q	7	1914	no previous epigastric pain	no definite finding	Hb 54% Rbc 2.4M Wbc 5500 Eos 2% Sed 1000	no	Peptic ulcer	no	Hyperacidity	Gastrectomy	definitely Ulcer (Long ulcers free ulcer)	249
1007	JOHN D. BROWN	47	Q	7	1914	no previous epigastric pain	no definite finding	Hb 54% Rbc 2.4M Wbc 5500 Eos 2% Sed 1000	no	Peptic ulcer	no	Hyperacidity	Gastrectomy	definitely Ulcer (Long ulcers free ulcer)	249
1008	JOHN D. BROWN	47	Q	7	1914	no previous epigastric pain	no definite finding	Hb 54% Rbc 2.4M Wbc 5500 Eos 2% Sed 1000	no	Peptic ulcer	no	Hyperacidity	Gastrectomy	definitely Ulcer (Long ulcers free ulcer)	249
1009	JOHN D. BROWN	47	Q	7	1914	no previous epigastric pain	no definite finding	Hb 54% Rbc 2.4M Wbc 5500 Eos 2% Sed 1000	no	Peptic ulcer	no	Hyperacidity	Gastrectomy	definitely Ulcer (Long ulcers free ulcer)	249
1010	JOHN D. BROWN	47	Q	7	1914	no previous epigastric pain	no definite finding	Hb 54% Rbc 2.4M Wbc 5500 Eos 2% Sed 1000	no	Peptic ulcer	no	Hyperacidity	Gastrectomy	definitely Ulcer (Long ulcers free ulcer)	249

CASE	NAME	AGE	SEX	MAR	OCCUPATION	PAST HISTORY	PRESENT HISTORY	PHYSICAL EXAM	LAB FINDINGS	PROVISIONAL DIAG	X-ray	GASTRIC ANALYSIS	OP. OPERATION	MICRO EXAM	REMARKS
1	WOMAN 54	54	F	1	HOUSEWIFE	NO	CC. Torso no pain no nausea no vomiting	no definite signs	Hb 41% Rbc 2.61 Wbc 6,200 Urine / spec	Carcinoma of Cardiac portion of Stomach	NO	gastric Low acidity	Gastric	NO	2400
2	WOMAN 50	50	F	1	HOUSEWIFE	NO	CC. Chronic regurgitation no vomiting	no definite signs	Hb 60% Rbc 5.22 Wbc 7,550 Urine / spec	Chronic gastritis Ulcer	NO	NO	Partial gastric ulcer in gastric ulcer	Chronic gastritis	2400
3	WOMAN 30	30	F	1	HOUSEWIFE	NO	CC. Chronic regurgitation no vomiting	no definite signs	Hb 36% Rbc 4.55 Wbc 8,600 Urine / spec	Marked anemia Abdominal pain in right epigastrium	Carcinoma of stomach in pyloric region & partial obli.	Low	Partial gastric ulcer	Adeno carcinoma	2400
4	WOMAN 30	30	F	1	HOUSEWIFE	NO	CC. Chronic regurgitation no vomiting	no definite signs	Hb 39% Rbc 2.10 Wbc 6,700 Urine / spec	epilepsy & hemorrhage	Duodenal ulcer	Moderate high acidity	Partial gastric ulcer in gastric region	Partial gastric ulcer	2400
5	WOMAN 30	30	F	1	HOUSEWIFE	NO	CC. Chronic regurgitation no vomiting	no definite signs	Hb 61% Rbc 2.65 Wbc 4,200 Urine / spec	Abdominal Tumor	Partial obli. gastric distension stomach	Low acidity	Partial gastric ulcer in gastric region	Partial gastric ulcer	2400
6	WOMAN 26	26	F	1	HOUSEWIFE	NO	CC. Chronic regurgitation no vomiting	no definite signs	Hb 38% Rbc 2.65 Wbc 4,200 Urine / spec	Abdominal Tumor	Tumor mass at epigastrium & middle stomach	Low	Partial gastric ulcer in gastric region	Adeno carcinoma	2400
7	WOMAN 21	21	F	1	HOUSEWIFE	NO	CC. Chronic regurgitation no vomiting	no definite signs	Hb 38% Rbc 4.22 Wbc 7,500 Urine / spec	Duodenal ulcer	NO	Hypochloric acid	Small gastric ulcer in gastric region	Duodenal ulcer	2400

CASE	NAME	AGE	SEX	RACE	DOB	PAST-HIST.	PRESENT H.	PHYSICAL EXAM	LAB.	Proctocolon Proc.	x-ray	Gastroc Analysis	Pp Operation	Microc copy	N.O.
1001	W. H. H. H.	47	F	F	W.H.	no	ce. chronic in 2nd & 3rd quadrants of the l.u.r.	fundamental epigastric region.	Hb 81% Pbc 88% Wbc 3800 V. normal	ch. duodenal and ch. l.u.r.	ch. duodenal and ch. l.u.r.	Hyper- acidity	Partial gastroctomy	no	2499
1002	W. H. H. H.	42	F	F	W.H.	no	ce. chronic in 2nd & 3rd quadrants of the l.u.r.	fundamental epigastric region.	Hb 81% Pbc 88% Wbc 3800 V. normal	ch. duodenal and ch. l.u.r.	ch. duodenal and ch. l.u.r.	Hyper- acidity	Partial gastroctomy + gastrojejunostomy	no	2499

เอกสารส่วนบุคคล ศาสตราจารย์นายแพทย์เสมอ พริงพวงแก้ว

สพ 1

..... 2/1:6 (ชื่อเรื่อง)
..... พ.ศ. 2494-2503 โรงพยาบาลศิริราช
.....

พ.ศ. 2494-2503

จำนวน 20 แผ่น

แฟ้มที่ 2

กล่องที่ 1

WOMEN'S AND CHILDREN'S HOSPITALS

MALIGNANCY IN CHILDREN

No.	H.N.	A.N.	Year.
1. Malignant Hepatoma & CA. bile duct.	8114	1306	1954
2. Neuroblastoma Rt. cheek.	15920	2761	1955
3. Nephroblastoma Lt. Kidney	16243	2902	1955
4. Wilms' tumour, Left.	27891	4837	1955
5. Malignant Hemangioendothelioma	52071	9390	1956
6. Papillary CA, of Testis, Right	62458	11394	1956
7. Wilms' Tumor, Left.	53053	9572	1956
8. Hypernephroma Left.	61772	11241	1956
9. Melanosarcoma	36637	6468	1956
10. Wilms' Tumor, Right.	67588	12406	1957
11. Metastatic Lymphosarcoma	92988	17121, 18929.	1957
12. Metastatic Teratoma.	89985	16499	1957
13. Teratoma of Soft Tissue, Left Renal Space	88670	16143	1957
14. Papillary Choroideum 7	85742	17622	1958
15. Malignant Hemangio-endothelioma	120667	22178	1958
16. Lymphosarcoma	117134	21561	1958
17. Lymphosarcoma, (intestine, mesentery)	121554	22321	1958
18. Round Cell Sarcoma Left Thigh	108061	19763	1958
19. Mediastinal Teratoma.	105714	19321	1958
20. Metastatic Teratoma.	-	16499	1958
21. Wilms' Tumour, Right.	115824	21365	1958
22. Adenocarcinoma of pancreas	134940	24501	1959
23. Retinoblastoma, Left Eye.	136998	24883	1959
24. Lymphosarcoma	142143	25696	1959
25. Lymphosarcoma	144619	26070	1959
26. Mesenteric Lymphosarcoma.	144082	25968	1959
27. Lymphosarcoma, Cervical Lymph Glands.	155452	27605	1959
28. Metastatic Tumor.	104940	29149	1959
29. Reticulum Cell Sarcoma, Rt. Cheek	201629	34381	1960
30. Mesenteric Lymphosarcoma.	178370	31775	1960
31. Mesenteric Lymphosarcoma	187096	-	1960
32. Neuroblastoma (Abdominal)	187718	32490	1960
33. Bronchogenic CA. Metastasis to Cardia and Pericardium.	173375	30729	1960
34. Lymphosarcoma, Mesenteric Lymph Glands.	188977	32676	1960
35. Lymphosarcoma.	191954	33178	1960
36. Wilms' Tumor, Right	197893	33871	1960
37. Wilms' Tumour	204980	34918	1960
38. Lymphosarcoma Mesenteric Lymph Glands.	205190	35009	1960
39. Neuroblastoma	189727	032893	1960
40. Melanoma Malignant.	211039	035913	1960

WOMEN'S AND CHILDREN'S HOSPITALS

MALIGNANCY IN CHILDREN

No.	H.N.	A.N.	Year.
1. Malignant Hepatoma & CA. bile duct.	8114	1306	1954
2. Neuroblastoma Rt. cheek.	15920	2761	1955
3. Nephroblastoma Lt. Kidney	16243	2902	1955
4. Wilms' tumour, Left.	27891	4837	1955
5. Malignant Hemangioendothelioma	52071	9390	1956
6. Papillary CA, of Testis, Right	62458	11394	1956
7. Wilms' Tumor, Left.	53053	9572	1956
8. Hypernephroma Left.	61772	11241	1956
9. Melanosarcoma	36637	6468	1956
10. Wilms' Tumor, Right.	67588	12406	1957
11. Metastatic Lymphosarcoma	92988	17121, 18929	1957
12. Metastatic Teratoma.	89985	16499	1957
13. Teratoma of Soft Tissue, Left Renal Space	88670	16143	1957
14. Papillary Choroidaem	85742	17622	1958
15. Malignant Hemangio-endotheliom	120667	22178	1958
16. Lymphosarcoma	117134	21561	1958
17. Lymphosarcoma, (intestine, mesentery)	121554	22321	1958
18. Round Cell Sarcoma Left Thigh	108061	19763	1958
19. Mediastinal Teratoma.	105714	19321	1958
20. Metastatic Teratoma.	-	16499	1958
21. Wilms' Tumour, Right.	115824	21365	1958
22. Adenocarcinoma of pancreas	134940	24501	1959
23. Retinoblastoma, Left Eye.	136998	24883	1959
24. Lymphosarcoma	142143	25696	1959
25. Lymphosarcoma	144619	26070	1959
26. Mesenteric Lymphosarcoma.	144082	25968	1959
27. Lymphosarcoma, Cervical Lymph Glands.	155452	27605	1959
28. Metastatic Tumor.	104940	29140	1959
29. Reticulum Cell Sarcoma, Rt. Cheek	201629	34381	1960
30. Mesenteric Lymphosarcoma.	178370	31775	1960
31. Mesenteric Lymphosarcoma	187096	-	1960
32. Neuriblastoma (Abdominal)	187718	32490	1960
33. Branchogenic CA. Metastasis to Cardia and Pericardium.	173375	30729	1960
34. Lymphosarcoma, Mesenteric Lymph Glands.	188977	32676	1960
35. Lymphosarcoma.	191954	33178	1960
36. Wilms' Tumor, Right	197893	33871	1960
37. Wilms' Tumour	204980	34918	1960
38. Lymphosarcoma Mesenteric Lymph Glands.	205190	35009	1960
39. Neuroblastoma	189727	032893	1960
40. Melanoma Malignant.	211039	035913	1960

MALIGNANCY IN CHILDREN
CHILDREN'S HOSPITAL, BANGKOK

I. Selected cases from those who had been treated with ENDOXAN - ASTA.

NO.	H.N.	A.N.	DATE	DIAGNOSIS
1.	188977	32676	July 6, 1960	Mesenteric Lymphosarc.
2.	173375 173639 173345	30329	July 4, 1960	Bronchogenic Carc.
3.	189727	032893	July 13, 1960	Neuroblastoma.
4.	205190	035009	Oct. 1960	Mesenteric Lymphosarc.
5.	197893	035229 033871	Sept. 1960	Metastatic Wilms'
6.	211039	035913	Dec. 23, 1960	Malignant Melanoma

7. A. case of Wilms' tumor which had been discharged recently during last month.
The patient's name is

H.N. 217496 A.N. 02962 DATE June 29, 1961

II. A case of Lymphosarcoma of mesenteric lymph gland that had been treated with Nitronin combined with surgery, but not ENDOXAN - ASTA.

9.	187096	032377	June 20, 1960	Mesent. Lymphosarc.
----	--------	--------	---------------	---------------------

((From the file of "Malignancy In Children" Kasem. Children's Hospital

WOMEN'S AND CHILDREN'S HOSPITALS

MALIGNANCY IN CHILDREN

No.	H.N.	A.N.	Year.
1. Malignant Hepatoma & CA. bile duct.	8114	1306	1954
2. Neuroblastoma Rt. cheek.	15920	2761	1955
3. Nephroblastoma Lt. Kidney	16243	2902	1955
4. Wilms' tumour, Left.	27891	4837	1955
5. Malignant Hemangioendothelioma	52071	9390	1956
6. Papillary CA, of Testis, Right.	62458	11394	1956
7. Wilms' Tumor, Left.	53053	9572	1956
8. Hypernephroma Left.	61772	11241	1956
9. Melanosarcoma	36637	6468	1956
10. Wilms' Tumor, Right.	67588	12406	1957
11. Metastatic Lymphosarcoma	92988	17121, 18929.	1957
12. Metastatic Teratoma.	89985	16499	1957
13. Teratoma of Soft Tissue, Left Renal Space.	88670	16143	1957
14. Papillary Choroidaem	85742	17622	1958
15. Malignant Hemangio-endothelioma	120667	22178	1958
16. Lymphosarcoma	117134	21561	1958
17. Lymphosarcoma, (intestine, mesentery)	121554	22321	1958
18. Round Cell Sarcoma Left Thigh	108061	19763	1958
19. Mediastinal Teratoma.	105714	19320	1958
20. Metastatic Teratoma.	-	16499	1958
21. Cerebral Tumour	123095	23092	1958
22. Brain Tumor	126421	22944	1958
23. Wilms' Tumour, Right.	115824	21365	1958
24. Adenocarcinoma of pancreas	134940	24501	1959
25. Retinoblastoma, Left Eye.	136998	24883	1959
26. Lymphosarcoma	142145	25696	1959
27. Lymphosarcoma	144619	26070	1959
28. Mesenteric Lymphosarcoma.	144082	25968	1959
29. Lymphosarcoma, Cervical Lymph Glands.	155452	27605	1959
30. Metastatic Tumor.	104940	29149	1959
31. Reticulum Cell Sarcoma, Rt. Cheek	201629	34381	1960
32. Mesenteric Lymphosarcoma.	178370	31775	1960
33. Mesenteric Lymphosarcoma	187096	-	1960
34. Neuroblastoma (Abdominal)	187718	32490	1960
35. Bronchogenic CA. Metastasis to Cardia and Pericardium.	173375	30729	1960
36. Lymphosarcoma, Mesenteric Lymph Glands.	188977	32676	1960
37. Lymphosarcoma.	191954	33178	1960
38. Wilms' Tumor, Right.	197893	33871	1960
39. Wilms' Tumour.	204980	34918	1960
40. Lymphosarcoma Mesenteric Lymph Glands.	205190	35009	1960
41. Neuroblastoma	189727	032893	1960
42. Melanoma Malignant.	211039	035913	1960

7 Years Period

Wilms' 6

Neuroblastoma 3

Melanoma 2

Lymphosarcoma 13

Krasem

Dec 12. 1961.

MALIGNANCY IN CHILDREN
CHILDREN'S HOSPITAL, BANGKOK.

I. Selected cases from those who had been treated with ENDOXAN - ASTA.

NO.	H.N.	A.N.	DATE	Diagnosis.
1.	189977	32676	July 6, 1960	Mesenteric Lymphosarc.
2.	173375 173639 173345	30329	July 4, 1960	Bronchogenic Carc.
3.	189727	032893	July 13, 1960	Neuroblastoma.
4.	205190	035009	Oct. 1960	Mesenteric Lymphosarc.
5.	197893	035229 033871	Sept. 1, 1960	Metastatic Wilms'
6.	211039	035913	Dec. 23, 1960	Malignant Melanoma

7. A case of Wilms' tumor which had been discharged recently during last month.

The patient's name is

HN. 217496. A.N. 02962. June 29, 1961.

II. A case of Lymphosarcoma of mesenteric lymph gland that had been treated with Nitronin combined with surgery, but not ENDOXAN-ASTA.

8.	187096	032377	June 20, 1960.	Mesent. Lymphosarc.
----	--------	--------	----------------	---------------------

(From the file of " Malignancy In Children "-----Kasem. Children's Hospital)

Aug. 10. 1961

Department of Radiology

2499 - 25

Carcinoma of cervix	448	cases.
Having complaint	39	cases.
X-Ray pelvis, lumbar spine	34	cases.
+ve. bony metastasis	7	cases.

= 1.56 %

total

20%

WOMEN'S AND CHILDREN'S HOSPITALS BANGKOK

SURGICAL DISEASES AND NUMBER OF CASES IN 1957
(Patients under twelve years of age)

1. Abscess	60
2. Abdominal pain, -ascariasis	8
3. Abdominal pain, -cause ?	8
4. Acute abdomen	3
5. Appendicitis, acute	8
6. Appendicitis, chronic	1
7. Anencephaly	2
8. Appendicial abscess	2
9. Abrasion	1
10. Brain abscess	1
11. Boils	1
12. Bleeding per vagina	8
13. Bleeding per rectum	3
14. Burns	11
15. Bleeding from gum	3
16. Bleeding from umbilicus	3
17. Cellulitis	59
18. Cleft lip, oblique facial cleft lip, left	1
19. Cooley's anemia	17
20. Cerebral concussion	7
21. Cyst, omental cyst	2
22. Cyst, pilonidal	1
23. Cyst, multiple cysts of chest	1
24. Cyst, mucous	1
25. Cyst, sebaceous	1
26. Cystocoele	1
27. Cyst, unknown nature	1
28. Cystic hygroma	1
29. Cystitis, chronic	1
30. Cleft palate	1
31. Contusion, cerebral	1
32. Contusion, abdominal	1
33. Contusion, labia majora	1
34. Concussion, abdominal	1
35. Deformity, flexion deformity of left hip joint joint	1
36. Dislocation, old, both hip joints	1
37. Dislocation, old, left hip joint	1
38. Dislocation, old , posterior dislocation, left hip joint	1
39. Duct, patent omphalomesenteric	1
40. Effusion, joints	2
41. Effusion, pleural	1
42. Epiphysitis	1
43. Epiphyseal seperation	1
44. Exomphalos	4
45. Fistula, rectovaginal	8
46. Fistula, urethrorectal	1
47. Fistula, rectoperineal	1
48. Furunculosis	3
49. F.B., in thigh	1
50. F.B., in right ear	1
51. F.B., in chest	1
52. Fracture, skull, compression	3
53. Fracture, skull, simple	1
54. Fracture, extremities, simple	13
55. Fracture, extremities, compound	1
56. Fracture, extremities, simple, incomplete	1
57. Fracture, extremities, simple, comminuted	1
58. Fracture, clavicle, simple	1
59. Fracture, ribs, simple	1
60. Granuloma	1

61. Gangrene right foot	1
62. Gangrene left leg	1
63. Gingivitis	1
64. Harelip, unilateral left	4
65. Harelip, unilateral left with cleft palate	4
66. Harelip, bilateral with cleft palate...	3
67. Harelip, with cleft palate	4
68. Harelip, unilateral, right with cleft palate	3
69. Hemorrhage, cerebral	4
70. Hemorrhage, intestinal	2
71. Hemorrhage, gastric	1
72. Hydrocephalus	15
73. Hernia, indirect inguinal, strangulated	0
74. Hernia, indirect inguinal, partial obstruction	2
75. Hernia, indirect inguinal, left, obstructed	2
76. Hernia, indirect inguinal, right, reducible	6
77. Hernia, indirect inguinal, left, reducible	6
78. Hernia, indirect inguinal	1
79. Hydrocoele, left	2
80. Hydrocoele, right	3
81. Hemangioma, infected	2
82. Hernia, umbilical	4
83. Hematoma	2
84. Imperforate anus	15
85. Intestinal obstruction, low gut	2
86. Intestinal obstruction, stenosis ileum & colon	1
87. Intestinal obstruction, narrowing of duodenum	1
88. Intestinal obstruction, intussusception ileo-ileal with ileo-cecal	1
89. Intestinal obstruction	5
90. Intestinal obstruction, ascariasis	1
91. Intestinal obstruction, volvulus terminal ileum	1
92. Intestinal obstruction, volvulus	1
93. Jaundice, obstructive; atresia of common bile duct	1
94. Kidney, atrophy of, right	1
95. Kidney, contusion of, left	1
96. Kidney, poly cystic	1
97. Mass, abdominal	4
98. Malformation, absence of parietal bones, nose, and scalp	1
99. Meningocele, lumbar	1
100. Meningocele, frontal	1
101. Meningocele, sacral	2
102. Megacolon	4
103. Microcephalus	2
104. Meningo-encephalocele	1
105. Molluscum contagiosum	1
106. Orchitis	1
107. Osteomyelitis, chronic	9
108. Omphalocele	1
109. Oesophagotracheal fistula	1
110. Peritonitis, localize	2
111. Peritonitis, generalize	4
112. Peritonitis, typhoid	1
113. Perforation, ileum	1
114. Perforation, mesenteric lymph glands	1
115. Perforation of liver, due to ascaris	1
116. Prolapse rectum	7
117. Polyp, rectal, single	12
118. Polyp, rectal, multiple	1
119. Phimosis	4
120. Paralysis right leg	2
121. Paralytic ileus	1
122. Pancreatitis, chronic	1
123. Pyloric stenosis, congenital hypertrophic	1
124. Pyloric stenosis, partial obstruction	1
125. Pylorospasm	1
126. Parotid abscess	2
127. Peptic ulcer	1
128. Periosteal defect, left radius	1
129. Renal calculi	1
130. Rupture of transverse colon	1

131.Ranula	2	
132.Retinoblastoma,R.E.	1	
133.Scald,secondary degree.....	12	
134.Scar,contracture,right arm with chest wall	1	
135.Scar,infected	1	
136.Stricture of rectum,P.O.....	1	
137.Spina bifida,sacral.....	1	
138.Sinus,pilonidal,sacrum.....	1	
139.Stomatitis	1	
140.Subluxation,right hip joint.....	1	
141.Tabes mesenterica.....	2	
142.Talipes equino varus.....	1	
143.T.B.Spine	10	
144.Thymus,enlarged	4	
145.Teratoma,left renal space.....	1	
146.Teratoma,mesenteric	1	
147.T.B.epiphysitis ,right femur.....	1	
148.T.B.distal end of femur,left.....	1	
149.T.B.hip,right	2	
150.T.B. hip,left	1	
151.Tumour,abdominal..METASTASIS LYMPHOSAR	1	COMA
152.Tear,first degree perineal.....	1	
153.Urethral calculus.....	3	
154.Ulcer	17	
155.Urachus,patent urachus.....	1	
156.Vesical calculus.....	9	
157.Vesical calculi	1	
158.Wound,gun shot	1	
159.Wound,stab	2	
160.Wound,infected	3	
161.Wound,traumatic wound of vagina,..	1	
162.Wound,traumatic wound of vulva,..	1	
163.Wound,tearing wound of palate.....	1	
164.Wound,infected wound of umbilicus.	1	
165.Wound,lacerated	2	
166.Wound,dog bite	1	
167.Wilms' tumour,right.....	1	

TOTAL

560

September 2,1958.

WOMEN'S AND CHILDREN'S HOSPITALS BANGKOK

GENERAL SURGICAL OPERATIONS

PROCEDURE IN 1957

1. Appendectomy	5
2. Appendectomy with O. & D.	1
3. Biopsy, liver	1
4. Bryant's traction	2
5. Bronchoscopy	1
6. Colostomy	5
7. Colostomy, permanent	3
8. Curettage	4
9. Debridement	4
10. Dilatation, rectum	2
11. Dislocation reduction	1
12. Excision	18
13. Excision, tract	3
14. Explore laparotomy	11
15. Eversion of sac	1
16. Extirpation of sac	2
17. Hernial reduction	1
18. Herniorrhaphy	5
19. Herniotomy	4
20. Ileocollectomy	1
21. Ileostomy	2
22. Intussusception reduction.....	1
23. Lysis, intestinal adhesion.....	1
24. Meatotomy	1
25. Nephrolithotomy	1
26. Open and drain(admission cases).....	70
27. Omental cystectomy	2
28. Open reduction, fracture.....	2
29. Perforation of anus, perineal.....	5
30. Plastic operations(harelip, meningocoele, etc.)	19
31. Plating	1
32. P.P.S. & Spica	20
33. Ramstedt's operation	1
34. Removal operations	10
35. Skin grafting, partial thickness.....	2
36. Skin grafting, whole thickness.....	1
37. Sequestrectomy	7
38. Skin traction	4
39. Splenectomy	6
40. Suprapubic cystotomy	9
41. Suture	8
42. Suture of liver	11
43. Suture of mesenteric lymph glands.....	1
44. Subaqueous drainage	1
45. Tracheotomy	1
46. Twisted removal	1

Total Operations

253

Συνοπτική Υδρονέφρωση, Wilms' Tumor + Neuroblastoma

W.W. 2494 - 2503

	Ευνοϊκή			Πονηρική		
	Υδρονέφρωση	Κακοήγητο Τumor of kidney	Neuroblastoma	Υδρονέφρωση	Wilms' Tumor	Neuroblastoma
W.W. 2494	-	1	-			
2495	+	-	-			
2496	1	-	-			
2497	-	-	-			
2498	1	1	-	-	1	-
2499	-	1	-	-	2	-
2500	-	-	-	-	-	-
2501	-	-	-	-	-	-
2502	1	-	-	-	-	-
2503	-	1 (Nephroblastoma)	-	1	3	1
Σύνολο	3	4	-	1	6	1

Συνοπτική με Σύντομη περιγραφή χαρακτηριστικών.

TUMOURS OF THE UTERUS

Malignant Tumours of the Uterus	2494	2495	2496	2497	2498	2499	2500	2501	2502	Total cases.
Carcinoma Cervix	10	22	46	43	79	58	107	73	66	504
Adeno - Acanthoma of Cervix	-	-	-	-	-	-	-	-	-	1
Condyloma of Cervix with } Carcinomatus change	-	-	-	-	3	1	-	-	-	4
Carcinoma Corpus	-	1	2	4	2	3	7	3	7	29
Sarcoma of Uterus	-	-	1	-	1	-	-	-	-	2
Chorio - epithelioma	1	-	3	4	1	1	5	2	1	18
Benign Tumours of the Uterus	2494	2495	2496	2497	2498	2499	2500	2501	2502	Total cases.
Myoma	22	32	17	37	45	40	45	46	25	319
Condyloma of Cervix	-	-	-	-	1	-	-	-	-	1
Cervical Polyps	5	11	8	10	-	2	15	19	7	77

TUMOURS OF THE UTERUS 955 CASES

2494 - 2502

MALIGNANT TUMOURS 558 cases = 58.42 %

1) Carcinoma Cervix	= 504 Cases = 52.4 %
2) Adeno - Acanthoma of Cervix	= 1 Case = 0.1 %
3) Condyloma of Cervix with Ca. Change	= 4 Cases = 0.4 %
4) Carcinoma Corpus	= 29 " = 2.2 %
5) Sarcoma of Uterus	= 2 " = 0.2 %
6) Chorio - epithelioma	= 18 " = 1.3 %

BENIGN TUMOURS 397 Cases = 41.58 %

1) Myoma	= 319 Cases = 33.3 %
2) Condyloma of Cervix	= 1 Case = 0.1 %
3) Cervical Polyps	= 77 Cases = 10 %

TUMOURS OF THE UTERUS

Malignant Tumours of the Uterus	2494	2495	2496	2497	2498	2499	2500	2501	2502	Total cases.
Carcinoma Cervix	10	22	46	43	79	58	107	73	66	504
Adeno - Acanthoma of Cervix	-	-	-	-	-	1	-	-	-	1
Condyloma of Cervix with } Carcinomatous change	-	-	-	-	3	1	-	-	-	4
Carcinoma Corpus	-	1	2	4	2	3	7	3	7	29
Sarcoma of Uterus	-	-	1	-	1	-	-	-	-	2
Chorio - epithelioma	1	-	3	4	1	1	5	2	1	18
Benign Tumours of the Uterus	2494	2495	2496	2497	2498	2499	2500	2501	2502	Total cases.
Myoma	22	32	17	37	45	40	45	46	25	319
Condyloma of Cervix	-	-	-	-	1	-	-	-	-	1
Cervical Polyps	5	11	8	10	-	2	15	19	7	77

TUMOURS OF THE UTERUS 955 CASES

2494 - 2502

MALIGNANT TUMOURS 558 cases = 58.42 %

1) Carcinoma Cervix	= 504 Cases = 52.4 %
2) Adeno - Acanthoma of Cervix	= 1 Case = 0.1 %
3) Condyloma of Cervix with Ca. Change	= 4 Cases = 0.4 %
4) Carcinoma Corpus	= 29 " = 2.2 %
5) Sarcoma of Uterus	= 2 " = 0.2 %
6) Chorio - epithelioma	= 18 " = 1.3 %

BENIGN TUMOURS 397 Cases = 41.58 %

1) Myoma	= 319 Cases = 33.3 %
2) Condyloma of Cervix	= 1 Case = 0.1 %
3) Cervical Polyps	= 77 Cases = 10 %

TUMOURS OF THE UTERUS

Malignant Tumours of the Uterus	2494	2495	2496	2497	2498	2499	2500	2501	2502	Total cases.
Carcinoma Cervix	10	22	46	43	79	58	107	73	66	504
Adeno - Acanthoma of Cervix	-	-	-	-	-	1	-	-	-	1
Condyloma of Cervix with } Carcinomatous change	-	-	-	-	3	1	-	-	-	4
Carcinoma Corpus	-	1	2	4	2	3	7	3	7	29
Sarcoma of Uterus	-	-	1	-	1	-	-	-	-	2
Chorio - epithelioma	1	-	3	4	1	1	5	2	1	18
Benign Tumours of the Uterus	2494	2495	2496	2497	2498	2499	2500	2501	2502	Total cases.
Myoma	22	32	17	37	45	40	45	46	25	319
Condyloma of Cervix	-	-	-	-	1	-	-	-	-	1
Cervical Polyps	5	11	8	10	-	2	15	19	7	77

TUMOURS OF THE UTERUS 955 CASES

2494 - 2502

MALIGNANT TUMOURS 558 cases = 58.42 %

1) Carcinoma Cervix	= 504 Cases = 52.4 %
2) Adeno - Acanthoma of Cervix	= 1 Case = 0.1 %
3) Condyloma of Cervix with Ca. Change	= 4 Cases = 0.4 %
4) Carcinoma Corpus	= 29 " = 2.2 %
5) Sarcoma of Uterus	= 2 " = 0.2 %
6) Chorio - epithelioma	= 18 " = 1.3 %

BENIGN TUMOURS 397 Cases = 41.58 %

1) Myoma	= 319 Cases = 33.3 %
2) Condyloma of Cervix	= 1 Case = 0.1 %
3) Cervical Polyps	= 77 Cases = 10 %

CARCINOMA CERVIX

MICROSCOPIC SECTION - GRADING OF TUMOUR

TOTAL = 566 CASES

- Adeno carcinoma of Cervix 5 cases.

- Grading of squamous cell carcinoma of cervix

TOTAL = 561 CASES

Border lined	=	12 cases.
Ca. in Situ	=	8 "
Grade 0 - 1	=	1 case
Grade 1	=	301 cases
Grade 1 - 2	=	76 "
Grade 2	=	154 "
Grade 2 - 3	=	8 "
Grade 3	=	None
Grade 3 - 4	=	1 case
Grade 5	=	None.

=====

CARCINOMA CERVIX

MICROSCOPIC SECTION - GRADING OF TUMOUR

TOTAL = 566 CASES

- Adeno carcinoma of Cervix 5 cases. = 0.9%

- Grading of squamous cell carcinoma of cervix = 99.1%

TOTAL = 561 CASES

Border lined	= 12 cases.	= 2.1%	} = 3.5%
Ca. in Situ	= 8 "	= 1.4%	
Grade 0 - 1	= 1 case	= 0.2%	} = 53.8%
Grade 1	= 301 cases	= 53.6%	
Grade 1 - 2	= 76 "	= 13.6%	} = 41.1%
Grade 2	= 154 "	= 27.5%	
Grade 2 - 3	= 8 "	= 1.4%	
Grade 3	= None		
Grade 3 - 4	= 1 case	= 0.2%	
Grade 5	= None.		

=====

CARCINOMA CERVIX

MICROSCOPIC SECTION - GRADING OF TUMOUR

TOTAL = 566 CASES

- Adeno carcinoma of Cervix 5 cases.

= 0.37 %

- Grading of squamous cell carcinoma of cervix

= 2.12 %

TOTAL = 561 CASES

Border lined	=	12 cases.
{ Ca. in Situ	=	8 "
{ Grade 0 - 1	=	1 case }
{ Grade 1	=	301 cases }
{ Grade 1 - 2	=	76 "
{ Grade 2	=	154 " }
{ Grade 2 - 3	=	8 "
Grade 3	=	None
Grade 3 - 4	=	1 case
Grade 5	=	None.

= 1.23 %

= 0.17 %

= 50.33 %

=====

CANCER AND NON CANCER TREATED WITH RADIUM AND DEEP X RAY.

NO.	SYSTEM	TOTAL	%	Radiation Induced CB
1.	Reproductive (except cervix)	105		① Skin Cancer ② Lung Cancer (Uranium miners)
2.	Cervix	352		③ Bone SARCOMA
3.	Alimentary	115		④ Thyroid CB (after Thyroid radiation)
4.	Breast	108		⑤ Pharyngeal CB after radiation for Kyntoxi cases or T.B. glands.
5.	Hemangioma	107		
6.	Lymphatic	69		
7.	Miscellaneous	46		
8.	Skin	56		
9.	Keloid	52		
	TOTAL	1010		

Spondylitis Therapy.

Thymus Therapy

Malformation

leukemia occurs 1 case in a year due
to radiation

Radiation 1:500 (Normal rate)

Double Mutation Rate to parents 1:485

Case of adenodiplosis - usual rate occurrence 1:9400
after radiation 1:5000

	2494	95	96	97	98	99	00	01	02	total
Ca. Corpus	0	1	3	4	3	3	7	3	6	30

Handwritten note: 2494-2504

	Total	Ratio	Age	Av. Age
Ca. Corpus:	24	1	34-78	54
Ca. Cervix	438	17.8	25-84	48 (40-50)

1. Age & Parity	Age	Av. Age	Nullipara	Premenopause
Novak&Kimbell	30-90	55	30 %	20 %- 30 %
Women's Hosp.	34-78	54	16.6%	20 %

Handwritten note: 100% Virgin - 100%

2. Late Menopause (way 1944)

Age of Menopause Ca corpus = 51.5 yrs, Normal = 46.5 yr

3. Obesity

4. Hypertension Bastinasse Over 150 mmHg. = 64 %
 Women's Hosp. - " - = 16.6 % (5/30)

5. Diabetes Mellitus

6. Oestrogenic Over Activity - No Method of Testing.

7. Polyps Way = 30 % Kollmier = 21 %

8. Fibroids Way = 30 % Women's Hosp. = 33 % (1/30)

9. Racial Factors Jeyes & Negro

10. Heredity *↓ Rare* *↑ Rare*

11. Previous Irradiat

12. Pathology

	Palmer 1079 Cases	Women's Hosp. 30 cases.
Adeno Carc	917 = 84.99 %	24 = 80 %
Adenosarcoma	121 = 11.21 %	1 = 3.3 %
Adenosarc & Ep. Cell metaplasia		1 = 3.3 %
Sarcoma of Muscle	31 = 2.96 %	2 = 6.8 %
Sarc. of Endometrium	3 = 0.28 %	1 = 3.3 %
Squamous cell Carc.	7 = 0.56 %	1 = 3.3 %
Cystic hyperplasia & Early Ca change		1 = 3.3 %

Handwritten note: Stage 2

Grade I
Grade II
Grade III
Grade IV

13. Histological Grading

14. Spread

15. Women's Hosp. Cervix 1 cases
Lungs 3 "
Ovaries 1 "
Parametrium 1 "
abd. viscerae 1 "

Generalized peritoneal spreading.

metastasis $\frac{7 \times 100}{30} = 23.3\%$

15. Classification

Editorial Committee

$\frac{1}{30} = 3.3$
 $\frac{22}{30}$
 $\frac{7}{30}$

Stage 0
Stage 1
" 2

Nearly Ca. Nature stage 0 100%
Confined in Uterus 100%
Outside Uterus 75%

Rickford

Group A
" B
" C
" D

Confined in End & uterine wall
To Cervical canal
To adnexae
To pelvic wall & distance -metastasis.

16. Symptoms & signs

17. Diagnosis ✓

18. Prognosis ✓

19. Treatment ✓

20. Recurrence.

Operation 100% - Women's Hospital

as 9 Generalized Ca. of Peritoneum
- no histological or Endometrial

.....

เอกสารส่วนบุคคล ศาสตราจารย์นายแพทย์เสม พริ้งพวงแก้ว

สบ 1

2/1.7 สืบค้นทางเอกสารก่อนกำหนด พ.ศ. 2561 (ชื่อเรื่อง)

2495-2499 โรงพยาบาลนครราชสีมา

พ.ศ. 2495 - 2499

จำนวน 4 แผ่น

แฟ้มที่ 2

กล่องที่ 1

Incidence of Eclampsia - Psychosis

Y N	จำนวนคน NS	จำนวน NS	Toxemia of pregnancy			Psychosis	N.B.
			Eclampsia	Pre- eclampsia	Total		
2495	3892	260	8	5	13 = 0.33%	1	คนที่นอนหลับ
2496	4802	484	12	18	30 = 0.7%	1	คนที่นอนหลับ
2497	6158	406	21	16	37 = 0.74%	- 1	คนที่นอนหลับ
2498	7331	626	34	31	65 = 0.88%	- 2	
2499	8617	792	22	34	61 = 0.78%	- 2	

3892
 4802

 8694
 6158

 14852
 7331

 22283
 8617

 31,900

8 cases

31,900 Total Anomalies

8 Cases

31,900 - 8

1
 1000

$\frac{8}{31,900} \times 1000$
 = $\frac{80}{319}$

319) 800 (2
 638

 162

31,900

$\frac{8}{31,900} \times 100$
 102

สถิติการคลอดลูกกำหนด ในโรงพยาบาลหญิง ตั้งแต่วันที่ ๑ มกราคม ถึง ๓๑ ธันวาคม ๒๔๕๕ รวม
 ๑ ปี คิดจากน้ำหนัก ๕๐๐ - ๒๒๐๐ กรัม มีจำนวน ๒๖๗ คน จากคนไข้มารับคลอดทั้งหมด ๔๖๑๗ คน คิดเป็นเปอร์เซ็นต์
 เท่ากับ ๓.๕ เปอร์เซ็นต์ และรับจากภายนอกโรงพยาบาล ๖๑ คน รวมเป็นทารกคลอดก่อนกำหนดที่แผนกทารก
 ในการรักษาพยาบาลตลอดปี ๓๒๘ คน

เดือน	จำนวนคนไข้มารับคลอด ทั้งหมด	เป็นทารกคลอดก่อน กำหนดที่คลอดใน โรงพยาบาล	เป็นทารกที่รับจากนอก โรงพยาบาล
มกราคม	๖๔๗ คน	๑๖ คน	๔ คน
กุมภาพันธ์	๖๒๒ คน	๑๒ คน	๑๐ คน
มีนาคม	๖๔๗ "	๒๕ "	๕ "
เมษายน	๖๔๔ "	๑๔ "	๗ "
พฤษภาคม	๖๖๐ "	๑๕ "	๕ "
มิถุนายน	๖๕๑ "	๒๓ "	๑ "
กรกฎาคม	๖๔๒ "	๓๒ "	๖ "
สิงหาคม	๗๖๗ "	๒๔ "	๒ "
กันยายน	๗๔๔ "	๓๖ "	๕ "
ตุลาคม	๘๖๔ "	๒๔ "	๔ "
พฤศจิกายน	๘๒๗ "	๒๔ "	๒ "
ธันวาคม	๗๕๘ "	๑๗ "	๒ "
รวมทั้งสิ้น	๘๖๑๗ คน	๒๖๗ คน	๖๑ คน

๑. รวมเด็กที่คลอดทั้งภายในและนอกโรงพยาบาล

๓๒๘ คน

๒. สาเหตุของการคลอดก่อนกำหนดมีดังนี้

๑. มารดาได้จับอันตราย - กระแทกกระแทก เช่น ตกบันได, รกเกาะแก่ง ๓๐ ราย
๒. มารดาเป็นโรคชักก่อนคลอด ๑๐ "
๓. มารดาเป็นไข้สูง ๓ "
๔. รกเกาะปากมดลูก ๔ "
๕. โรคพิษแห่งครรภ์ ๑๐ "
๖. ไข้มารับ ๓ "

๗. บิด	๘. ราย
๘. กามโรค	๑๕ ราย
๙. โรคหัวใจ	๒ "
๑๐. แผล ๓ คน	๑ "
๑๑. แผล ๒ คน	๒๒ "
รวมทั้งสิ้น	๑๑๖ ราย

๓. แผล ๓ คน	- คนแรกหนัก	๑๗๐๐ กรัม
	- คนที่ ๒ หนัก	๕๐๐ "
	- คนที่ ๓ หนัก	๑๖๒๐ "

- เด็กแรก ๓ คนนี้ คนที่ ๑ - ๒ ถึงแก่กรรม เนื่องจากท้องเต็ม
คนที่ ๓ เสียใจ ๑ เดือนจนน้ำหนักได้ ๒๐๐๐ กรัม

แข็งแรงจนย้ายกลับบ้าน.

๔. เป็นทารกคลอดก่อนกำหนดไม่ทราบสาเหตุ ๒๑๒ ราย

๕. สถิติการตายและการมีชีวิตอยู่ของเด็ก ซึ่งมีน้ำหนักต่าง ๆ กัน ดังนี้

๕๐๐ - ๑๐๐๐ กรัม	๑๐๐๐ - ๑๕๐๐ กรัม	๑๕๐๐ - ๒๐๐๐ กรัม	๒๐๐๐ - ๒๕๐๐ กรัม
เกิดทั้งหมด ๒๔ คน	เกิดทั้งหมด ๕๔ คน มีชีวิต ๒๑ คน	เกิดทั้งหมด ๑๐๕ คน มีชีวิต ๗๖ คน	เกิดทั้งหมด ๑๑๗ คน มีชีวิต ๕๕ คน
ถึงแก่กรรมทั้งหมด ทั้งนี้	ถึงแก่กรรม ๓๓ คน	ถึงแก่กรรม ๕๓ คน ถึงแก่กรรมภายใน ๒๔ ชม. ๒๔ คน	ถึงแก่กรรม ๒๒ คน ถึงแก่กรรมภายใน ๒๔ ชม. ๑๒ คน
ถึงแก่กรรมภายใน ๔๕ ชม. ๒๐ คน	ถึงแก่กรรมภายใน ๒๔ ชม. ๑๔ คน	ถึงแก่กรรมภายใน ๑ สัปดาห์ ๑๕ คน ถึงแก่กรรมภายใน ๒ สัปดาห์ ๕ คน	ถึงแก่กรรมภายใน ๓ วัน ๕ คน ถึงแก่กรรมภายใน ๑๕ วัน ๕ คน
ถึงแก่กรรมภายใน ๓ วัน ๖ คน	ถึงแก่กรรมภายใน ๑ สัปดาห์ ๗ คน	ถึงแก่กรรมภายใน ๑ เดือน ๖ คน	เสียชีวิต ๗๖%
เลี้ยงจนอายุ ๑ เดือน แล้วถึงแก่กรรม ๒ คน	ถึงแก่กรรมภายใน ๑ สัปดาห์ ๑๕ คน	ถึงแก่กรรมภายใน ๑ เดือน ๖ คน	
เสียชีวิต ๐ %	ถึงแก่กรรมภายใน ๑ เดือน ๔ คน	ถึงแก่กรรมภายใน ๒ สัปดาห์ ๑๕ คน	
	เสียชีวิต ๓๕ %	เสียชีวิต ๓๕ %	

๖. สาเหตุของการตายของทารกคลอดก่อนกำหนด

๑. Anoxaemia เนื่องจาก Caesarian Section ๖ ราย
๒. Birth injuries จากการ forcep ๒ ราย

๒. หนองเยื่อ มี	Mucous	อุกขวางทางเดินหายใจ	๓๘	ราย
๔. น้ำตาเหนียวมี	Sclerema		๑๗	"
๕. เด็ก Toxic	หลุดจากมรดาเป็น	Eclampsia	๑๑	"
๖. หัวเหลืองจืด			๑๒	"
๗. Gastrointestinal Disturbances			๓๕	"
๘. Birth injuries	Prolong Second Stage		๑๕	"
	เนื่องจาก			

รวมทั้งสิ้น ๑๓๖ ราย

๗. สรุปปี ๒๔๙๙

แผนกทารกคลอดก่อนกำหนดครบคนไข		๓๒๘ คน
ถึงแก่กรรม		๑๓๖ คน
เกิดเป็นเปอร์เซ็นต์โดยงรอด		— ๖๐ %
" ถึงแก่กรรม		— ๔๐ %
เฉลี่ยตามน้ำหนักเด็กงรอดเกิดเป็นเปอร์เซ็นต์ดังนี้		
น้ำหนัก ๕๐๐-๑๐๐๐ กรัม	เด็กงรอด	๐%
" ๑๐๐๐ - ๑๕๐๐ กรัม	เด็กงรอด	๓๕%
" ๑๕๐๐ - ๒๐๐๐ กรัม	เด็กงรอด	๓๕%
" ๒๐๐๐ - ๒๕๐๐ กรัม	เด็กงรอด	๓๕%

กระดานข่าว

ผู้ร่าง

วันที่

ผู้ตรวจ

วันที่

เรื่อง x-ray Exam in Pregnancy ^{การตั้งครรภ์}

1. Verification of multiple Pregnancy
 2. Mal-presentation.
 3. Fetal abnormalities
 4. Hydramnios.
 5. Intra-uterine Death.
 6. Placenta Previa
 7. Hydatidiform mole
 8. Bony abnormalities
- Only in Primigravida
Suspected disproportion.
Breech presentation.

Pelvimetry

Mass ~~x~~-ray Chest Survey

4.7 millions examination
17,853 Pulm. T.B.
63,000 other abnormalities
2362 Lung CA.
12,000 Cardiac abnormalities
Could have produced at most 20 cases of leukemia.

X-ray Hazard

13,382 treated for gonorrhea by x-ray.
Leukemia occur in 38 cases. Causal rate
is about 2%

[13]

Thyroxine - 5-7 mc.
Thyroid CA. 100-150 mc

Beware of the
disposal of patient
Excreted
in urine used for
administration.

เอกสารส่วนบุคคล ศาสตราจารย์นายแพทย์เสม พริ้งพวงแก้ว

สพ 1

• 2/1.8 รูปสเก็ตของหม่อมหลวงพิริย (ชื่อเรื่อง)
พ.ศ. 2494 - 2503 โรงพยาบาลสงขลา

พ.ศ. 2494 - 2503


จำนวน 20 แผ่น

แฟ้มที่ 2

กล่องที่ 1

STATISTICAL SUMMARY OF OBSTETRICAL DEPARTMENT
IN WOMEN'S HOSPITAL

The total number of deliveries (2494 - 2503)



Multiple pregnancy

Twin	I: 109
Triplet	I: 8,742
Quadruplet	I: 34,069
Siamese twin	I: 34,069

Australia
1:80

Method of delivery

Normal deliveries	91.20	%
Abnormal deliveries	8.80	%
B.B.A. (305 cases)	0.31	%

The total number of live birth

Full term	77,378	=	95.79	%
Premature	3,047	=	4.21	%

13.4%

Sex

Male	52.26	%
Female	47.74	%

Weight

3,102 gm.

Length

49.77 cm.

Weight of placenta

579 gm.

Length of umbilical Cord

49.77 cm.

Fetal skull

	<u>Diameter</u>	<u>Circumference</u>
M.O.	13.5 cm.	37.5 cm.
P.O.	11.0 "	34.0 "
S.O.D.	9.5 "	31.5 "
B.P.	9.2 "	
B.T.	7.0 "	

PelvisAustralia

Interspinous	24	cm.
Intracristal	26	"
External conjugate	18	"

Position and presentation

O.L.A.	62.9	%
O.R.A.	33.81	%
O.P.P.	0.80	%
O.L.P.	0.43	%
O.R.P.	0.18	%
Breech	1.7	%
Transverse	0.19	%
Face	0.08	%
Brow	0.00001	%

Congenital anomalies

1. Abnormality of extremities	37.9	%
2. " " of G.I. tract	26.1	%
3. " " of nervous system	20.3	%
4. " " of urogenital system	5.3	%
5. " " of E.E.N.T.	2.1	%
6. Miscellaneous	8.3	%

1%

Blood Kahn

	<u>2494</u>	<u>2498</u>	<u>2502</u>	<u>2503</u>
3+	-	-	0.62 %	0.86 %
4+	7.8 %	7.18 %	1.5 %	2.47 %

Operative obstetric deliveries

Forceps extraction	2.49	%
Caesarean section	2.08	%
Craniotomy	0.04	%
Evisceration & Embryotomy	0.01	%
Internal version	0.23	%
Manual removal of placenta	0.70	%

12%

3-7%

Postpartum haemorrhage

Incidence	2.05	%
Grade I (300-500)	96.4	%
Grade II (500-1000)	37.4	%

→ 3.3%

(20 minutes
less 64)

Grade	III	(1000-1500)	5.8	%
Grade	IV	(1500-2000)	0.4	%

The Torsion of Pregnancy

Incidence	I:153	%
Variety		
1. Pre-eclampsia	76.28	%
2. Eclampsia	23.72	%
a. Antepartum	19.03	%
b. Postpartum	4.69	%
Parity		
Primi = I:1.4 = Multigravidarum		
Maternal mortality rate	3.85	%
(7 cases undeliveries; 1 case B.B.A.)		
Fetal mortality rate	9.25	%
Neonatal death	1.28	%
Intrauterine death	7.96	%

Pre-eclap = 3%

0.2%

Accidental hemorrhage

Incidence	I:503	
Primigravidarum	6.9	%
Multigravidarum	92.5	%
Maternal mortality rate	2.1	%
Fetal mortality rate	68.1	%

= 1.4%

Placenta previa

Incidence	I:206 (0.5%)	
Primigravidarum	11.05	%
Multigravidarum	88.93	%
Variety		
- total	39.9	%
- partial	38.3	%
- low insertion	22.2	%
Maternal mortality rate	2.54	%
Fetal mortality rate	40.45	%

→ 0.7%

Rupture uteri

Incidence

1:1,198

Primigravidarum

10.4 %

Multigravidarum

98.7 %

Causes:

A. Spontaneous rupture

a. previous section scar

17.45 %

b. other causes

72.05 %

B. Obstetrical trauma

21 %

Maternal mortality rate

14.42 %

Fetal mortality rate

85.92 %

Maternal mortality rate (~~1~~^{For} 1,000 live birth)

1. Due to obstetrical causes = 0.52:1,000 live birth

2. Due to associated diseases = 0.39:1,000 live birth

3. Due to toxic condition before admission = 0.34:1,000 live birth

Fetal mortality rate

In delivery room = 20.21:1,000 live birth

I. Intrauterine death = 15.6 :1,000 live birth

Neonatal death - full term = 2.59:1,000 live birth


- premature = 2.02:1,000 live birth

Neonatal Death
= 12/1000

Stillbirth
12/1000

STATISTICAL SUMMARY OF OBSTETRICAL DEPARTMENT
IN WOMEN'S HOSPITAL

The total number of deliveries (2494-2503)



<u>Multiple pregnancy</u>			<u>Method of deliveries</u>		
Twin	1:109		Normal deliveries	91.20	%
Triplet	1:8,724		Abnormal deliveries	8.80	%
Quadrupet	1:34,069		B.B.A. (305) cases)	0.31	%
Siamese twin	1:34,069		<u>The total number of live birth</u>		
<u>Sex</u>					
Male	52.26	%	Full term	77,378 =	95.79 %
Female	47.74	%	Premature	3,047 =	4.21 %
<u>Weight</u>		<u>Length</u>	<u>Congenital anomalies</u>		
3,102 gm.		49.77 cm.	1. Abnormality of extremities	37.9	%
<u>Weight of placenta</u>		<u>Length of umbilicus</u>	2. " " of G.I. tract	26.1	%
579 gm.		49.77 cm.	3. " " of nervous system	20.3	%
<u>Fetal skull</u>			4. " " of urogenital system	5.3	%
<u>Diameter</u>		<u>Circumference</u>	5. " " of E.E.N.T.	2.1	%
M.O. 13.5 cm.		37.5 cm.	6. Miscellaneous	8.3	%
F.O. 11.0 "		34.0 "	<u>Position and presentation</u>		
S.O.B. 9.5 "		31.5 "	O.L.A.	62.9	%
B.P. 9.2 "			O.R.A.	33.81	%
B.T. 7.0 "			O.P.P.	0.80	%
<u>Pelvis</u>			O.L.P.	0.43	%
Interspinous	24	cm.	O.R.P.	0.18	%
Intercristal	26	"	Breech	1.7	%
External conjugate	18	"	Transverse	0.19	%
<u>Blood Kahn</u>			Face	0.08	%
2494	2498	2502 2503	Brow	0.0001	%
3 ⁺	-	0.62 % 0.86 %			
4 ⁺	7.8 %	7.18 % 1.5 % 2.47 %			

Operative obstetric deliveries

Forceps extraction	2.49	%
Caesarean section	2.08	%
Craniotomy	0.04	%
Evisceration & Embryotomy	0.01	%
Internal version	0.23	%
Manual removal of placenta	0.07	%

Postpartum haemorrhage

Incidence	12.05	%
Grade I (300-500)	56.4	%
Grade II (500-1000)	37.4	%
Grade III (500-1500)	5.8	%
Grade IV (500-2000)	0.4	%

The Toxemia of Pregnancy

Incidence	1:153	
Variety		
1. Pre-eclampsia	76.28	%
2. Eclampsia	23.72	%
a. Antepartum	19.03	%
b. Postpartum	4.69	%

Parity

Primi = 1:1.4 = Multigravidarum

Maternal mortality rate	3.85	%
(7 cases undelivered, 1 B.B.A.)		
Fetal mortality rate	9.25	%
Neonatal death	1.28	%
Intrauterine death	7.96	%

Maternal mortality rate
(in 1,000 live birth)

1. Due to obstetrical causes	=	0.52:1,000	live birth
2. Due to associated causes	=	0.39:1,000	live birth
3. Due to condition before admission	=	0.34:1,000	live birth

Fetal mortality rate

In delivery room	=	20.21: 1,000	live birth
1. Intrauterine death		15.6:1,000	live birth
2. Neonatal death			
- full term		2.59:1,000	live birth
- premature		2.02:1,000	live birth

Placenta previa

Incidence	1:206	%
Primigravidarum	11.05	%
Multigravidarum	88.93	%
Variety		
- total	39.9	%
- partial	38.3	%
- low insertion	22.2	%

Maternal mortality rate	2.54	%
Fetal mortality rate	40.45	%

Accidental haemorrhage

Incidence	1:503	
Primigravidarum	6.9	%
Multigravidarum	92.5	%
Maternal mortality rate	2.1	%
Fetal mortality rate	68.1	%

Ruptured uteri

Incidence	1:1,198	
Primigravidarum	10.4	%
Multigravidarum	98.7	%

Causes:

A. Spontaneous rupture


a. previous section scar	17.45	%
b. other causes	72.05	%

B. Obstetrical trauma 21 %

Maternal mortality rate	14.42	%
Fetal mortality rate	85.92	%

STATISTICAL SUMMARY OF OBSTETRICAL DEPARTMENT

IN WOMEN'S HOSPITAL

2.The total number of deliveries (2494-2503)

<u>Multiple pregnancy</u>		<u>Method of deliveries</u>	
Twin	1:109	Normal deliveries	91.20 %
Triplet	1:8,724	Abnormal deliveries	8.80 %
Quadrupet	1:34,069	B.B.A. (305) cases)	0.31 %
Siamese twin	1:34,069	<u>The total number of live birth</u>	

<u>Sex</u>		<u>Sex</u>	
Male	52.26 %	Full term	77,378 = 95.79 %
Female	47.74 %	Premature	3,047 = 4.21 %

<u>Weight</u>		<u>Length</u>		<u>Congenital anomalies</u>	
3,102 gm.	49.77 cm.	1.Abnormality of extremities	37.9 %	2. " " of G.I. tract	26.1 %
<u>Weight of placenta</u>		<u>Length of umbilicus</u>		3. " " of nervous system	20.3 %
579 gm.	49.77 cm.	4. " " of urogenital system	5.3 %	5. " " of E.E.N.T.	2.1 %
<u>Fetal skull</u>		6. Miscellaneous	8.3 %	<u>Position and presentation</u>	
<u>Diameter</u>		<u>Circumference</u>		O.L.A.	62.9 %
M.O. 13.5 cm.	37.5 cm.			O.R.A.	33.81 %
F.O. 11.0 "	34.0 "			O.P.P.	0.80 %
S.O.B. 9.5 "	31.5 "			O.L.P.	0.43 %
B.P. 9.2 "				O.R.P.	0.18 %
B.T. 7.0 "				Breech	1.7 %

B.P.	9.2	"	O.L.A.	62.9	%		
B.T.	7.0	"	O.R.A.	33.81	%		
<u>Pelvis</u>			O.P.P.	0.80	%		
Interspinous	24	cm.	O.L.P.	0.43	%		
Intercristal	26	"	O.R.P.	0.18	%		
External conjugate	18	"	Breech	1.7	%		
<u>Blood Kahn</u>			Transverse	0.19	%		
	2494	2498	2502	2503	Face	0.08	%
3 ⁺	-	-	0.62 %	0.86 %	Brow	0.0001	%
4 ⁺	7.8 %	7.18 %	1.5 %	2.47 %			

Operative obstetric deliveries

Forceps extraction	2.49	%
Caesarean section	2.08	%
Craniotomy	0.04	%
Evisceration & Embryotomy	0.01	%
Internal version	0.23	%
Manual removal of placenta	0.07	%

Postpartum haemorrhage

Incidence	121.05	%
Grade I (300-500)	56.4	%
Grade II (500-1000)	37.4	%
Grade III (500-1500)	5.8	%
Grade IV (500-2000)	0.4	%

The Toxemia of Pregnancy

Incidence	1:153
Variety	
1. Pre-eclampsia	76.28 %
2. Eclampsia	23.72 %
a. Antepartum	19.03 %
b. Postpartum	4.69 %

Parity

Primi = 1:1.4 = Multigravidarum

Maternal mortality rate	3.85	%
(7 cases undelivered, 1 B.B.A.)		
Fetal mortality rate	9.25	%
Neonatal death	1.28	%
Intrauterine death	7.96	%

Maternal mortality rate
(in 1,000 live birth)

1. Due to obstetrical causes	=	0.52:1,000	live birth
2. Due to associated causes	=	0.39:1,000	live birth
3. Due to condition before admission	=	0.34:1,000	live birth

Fetal mortality rate

In delivery room	=	20.21: 1,000	live birth
1. Intrauterine death		15.6:1,000	live birth
2. Neonatal death			
- full term		2.59:1,000	live birth
- premature		2.02:1,000	live birth

Placenta previa

Incidence	1:206	%
Primigravidarum	11.05	%
Multigravidarum	88.93	%
Variety		
- total	39.9	%
- partial	38.3	%
- low insertion	22.2	%

Maternal mortality rate	2.54	%
Fetal mortality rate	40.45	%

Accidental haemorrhage

Incidence	1:503	
Primigravidarum	6.9	%
Multigravidarum	92.5	%
Maternal mortality rate	2.1	%
Fetal mortality rate	68.1	%

Ruptured uteri

Incidence	1:1,198	
Primigravidarum	10.4	%
Multigravidarum	98.7	%

Causes:

A. Spontaneous rupture


a. previous section scar	17.45	%
b. other causes	72.05	%

B. Obstetrical trauma

Maternal mortality rate	14.42	%
Fetal mortality rate	85.92	%

STATISTICAL SUMMARY OF OBSTETRICAL DEPARTMENT
IN WOMEN'S HOSPITAL

The total number of deliveries (2494-2503)



Twin	1:109	Normal deliveries	91.20	%
Triplet	1:8,724	Abnormal deliveries	8.80	%
Quadrupet	1:34,069	B.B.A. (305) cases)	0.31	%
Siamese twin	1:34,069	<u>The total number of live birth</u>		

Sex

Male	52.26	%	Full term	77,378	=	95.79	%
Female	47.74	%	Premature	3,047	=	4.21	%

Weight

Length

Congenital anomalies

3,102	gm.	49.77	cm.	1. Abnormality of extremities	37.9	%	
<u>Weight of placenta</u> <u>Length of umbilicus</u>				2. " " of G.I. tract	26.1	%	
579	gm.	49.77	cm.	3. " " of nervous system	20.3	%	
<u>Fetal skull</u>				4. " " of urogenital system	5.3	%	
<u>Diameter</u> <u>Circumference</u>				5. " " of E.E.N.T.	2.1	%	
M.O.	13.5	cm.	37.5	cm.	6. Miscellaneous	8.3	%

Position and presentation

				O.L.A.	62.9	%
				O.R.A.	33.81	%
				O.P.P.	0.80	%
				O.L.P.	0.43	%
				O.R.P.	0.18	%
				Breech	1.7	%
				Transverse	0.19	%
				Face	0.08	%
				Brow	0.0001	%

Blood Kahn

	2494	2498	2502	2503
3+	-	-	0.62 %	0.86 %
4+	7.8 %	7.18 %	1.5 %	2.47 %

Operative obstetric deliveries

Forceps extraction	2.49	%
Caesarean section	2.08	%
Craniotomy	0.04	%
Evisceration & Embryotomy	0.01	%
Internal version	0.23	%
Manual removal of placenta	0.07	%

Postpartum haemorrhage

Incidence	12.05	%
Grade I (300-500)	56.4	%
Grade II (500-1000)	37.4	%
Grade III (500-1500)	5.8	%
Grade IV (500-2000)	0.4	%

The Toxemia of Pregnancy

Incidence	1:153
Variety	
1. Pre-eclampsia	76.28 %
2. Eclampsia	23.72 %
a. Antepartum	19.03 %
b. Postpartum	4.69 %

Parity

Primi = 1:1.4 = Multigravidarum

Maternal mortality rate	3.85	%
(7 cases undelivered, 1 B.B.A.)		
Fetal mortality rate	9.25	%
Neonatal death	1.28	%
Intrauterine death	7.96	%

Maternal mortality rate
(in 1,000 live birth)

1. Due to obstetrical causes	=	0.52:1,000	live birth
2. Due to associated causes	=	0.39:1,000	live birth
3. Due to condition before admission	=	0.34:1,000	live birth

Fetal mortality rate

In delivery room	=	20.21: 1,000	live birth
1. Intrauterine death		15.6:1,000	live birth
2. Neonatal death			
- full term		2.59:1,000	live birth
- premature		2.02:1,000	live birth

Placenta previa

Incidence	1:206	%
Primigravidarum	11.05	%
Multigravidarum	88.93	%
Variety		
- total	39.9	%
- partial	38.3	%
- low insertion	22.2	%

Maternal mortality rate	2.54	%
Fetal mortality rate	40.45	%

Accidental haemorrhage

Incidence	1:503	
Primigravidarum	6.9	%
Multigravidarum	92.5	%
Maternal mortality rate	2.1	%
Fetal mortality rate	68.1	%

Ruptured uteri

Incidence	1:1,198	
Primigravidarum	10.4	%
Multigravidarum	98.7	%

Causes:

A. Spontaneous rupture


a. previous section scar	17.45	%
b. other causes	72.05	%

B. Obstetrical trauma

Maternal mortality rate	14.42	%
Fetal mortality rate	85.92	%

STATISTICAL SUMMARY OF OBSTETRICAL DEPARTMENT
IN WOMEN'S HOSPITAL

The total number of deliveries (2494-2503)



Twin	1:109	Normal deliveries	91.20	%
Triplet	1:8,724	Abnormal deliveries	8.80	%
Quadrupet	1:34,069	B.B.A. (305) cases)	0.31	%
Sianese twin	1:34,069	<u>The total number of live birth</u>		

Sex

Male	52.26	%	Full term	77,378	=	95.79	%
Female	47.74	%	Premature	3,047	=	4.21	%

Weight

Length

Congenital anomalies

3,102	gm.	49.77	cm.	1. Abnormality of extremities	37.9	%	
<u>Weight of placenta</u> <u>Length of umbilicus</u>				2. " " of G.I. tract	26.1	%	
579	gm.	49.77	cm.	3. " " of nervous system	20.3	%	
<u>Fetal skull</u>				4. " " of urogenital system	5.3	%	
<u>Diameter</u> <u>Circumference</u>				5. " " of E.E.N.T.	2.1	%	
M.O.	13.5	cm.	37.5	cm.	6. Miscellaneous	8.3	%

Position and presentation

				O.L.A.	62.9	%
				O.R.A.	33.81	%
				O.P.P.	0.80	%
				O.L.P.	0.43	%
				O.R.P.	0.18	%
				Breech	1.7	%
				Transverse	0.19	%
				Face	0.08	%
				Brow	0.0001	%

Blood Kahn

	2494	2498	2502	2503
3+	-	-	0.62 %	0.86 %
4+	7.8 %	7.18 %	1.5 %	2.47 %

Operative obstetric deliveries

Forceps extraction	2.49	%
Caesarean section	2.08	%
Craniotomy	0.04	%
Evisceration & Embryotomy	0.01	%
Internal version	0.23	%
Manual removal of placenta	0.07	%

Postpartum haemorrhage

Incidence	12.05	%
Grade I (300-500)	56.4	%
Grade II (500-1000)	37.4	%
Grade III (500-1500)	5.8	%
Grade IV (500-2000)	0.4	%

The Toxemia of Pregnancy

Incidence	1:153	
Variety		
1. Pre-eclampsia	76.28	%
2. Eclampsia	23.72	%
a. Antepartum	19.03	%
b. Postpartum	4.69	%

Parity

Primi = 1:1.4 = Multigravidarum

Maternal mortality rate	3.85	%
(7 cases undelivered, 1 B.B.A.)		
Fetal mortality rate	9.25	%
Neonatal death	1.28	%
Intrauterine death	7.96	%

Placenta previa

Incidence	1:206	%
Primigravidarum	11.05	%
Multigravidarum	88.93	%
Variety		
- total	39.9	%
- partial	38.3	%
- low insertion	22.2	%

Maternal mortality rate	2.54	%
Fetal mortality rate	40.45	%

Accidental haemorrhage

Incidence	1:503	
Primigravidarum	6.9	%
Multigravidarum	92.5	%
Maternal mortality rate	2.1	%
Fetal mortality rate	68.1	%

Ruptured uterus

Incidence	1:1,198	
Primigravidarum	10.4	%
Multigravidarum	98.7	%

Causes:

A. Spontaneous rupture

a. previous section scar	17.45	%
b. other causes	72.05	%

B. Obstetrical trauma

Maternal mortality rate	14.42	%
Fetal mortality rate	85.92	%

Maternal mortality rate
(in 1,000 live birth)

1. Due to obstetrical causes	= 0.52:1,000	live birth
2. Due to associated causes	= 0.39:1,000	live birth
3. Due to condition before admission	= 0.34:1,000	live birth

Fetal mortality rate

In delivery room

1. Intrauterine death
2. Neonatal death


- full term

- premature

= 20.21: 1,000	live birth
15.6:1,000	live birth
2.59:1,000	live birth
2.02:1,000	live birth

STATISTICAL SUMMARY OF OBSTETRICAL DEPARTMENT
IN WOMEN'S HOSPITAL

The total number of deliveries (2494 - 2503)



<u>Multiple pregnancy</u>		<u>Method of deliveries</u>	
Twin	1:109	Normal deliveries	91.20 %
Tripet	1:8,724	Abnormal deliveries	8.80 %
Quadrupet	1:34,069	B.B.A. (305 cases)	0.31 %
Siamese twin	1:34,069		

<u>Sex</u>		<u>The total number of live birth</u>	
Male	52.26 %	Full term	77,378 = 95.79 %
Female	47.74 %	Premature	3,047 = 4.21 %

<u>Weight</u>	<u>Length</u>	<u>Congenital anomalies</u>	
3,102 gm.	49.77 cm.	1. Abnormality of extremities	37.9 %
<u>Weight of placenta</u> <u>length of umbilicus</u>		2. " " of G.I. tract	26.1 %
579 gm.	49.77 cm.	3. " " of nervous system	20.3 %
<u>Fetal skull</u>		4. " " of urogenital system	5.3 %
<u>Diameter</u>	<u>Circumference</u>	5. " " of E.E.N.T.	2.1 %
M.O. 13.5 cm.	37.5 cm.	6. Miscellaneous	8.3 %
F.O. 11.0 "	34.0 "		

<u>Position and presentation</u>	
S.O.B. 9.5 "	31.5 "
B.P. 9.2 "	
B.T. 7.0 "	
O.L.A.	62.9 %
O.R.A.	33.81 %
O.P.P.	0.80 %
O.L.P.	0.43 %
O.R.P.	0.18 %
Breech	1.7 %
Transverse Face	0.19 %
	0.08 %

<u>Blood Kahn</u>			
	2494	2498	2502 2503
3	-	-	0.62% 0.86%
4	7.8%	7.18%	1.5 % 2.47%

Brow

Operative obstetric deliveries

Forceps extraction	2.49	%
Caesarean section	2.08	%
Craniotomy	0.04	%
Evisceration and Embryotomy	0.01	%
Internal version	0.23	%
Manual removal of placenta	0.70	%

Postpartum haemorrhage

Incidence	2.05	
Grade I (300-500)	56.4	%
Grade II (500-1000)	37.4	%
Grade III (1000-1500)	5.8	%
Grade IV (1500-2000)	0.4	%

The Toxemia of Pregnancy

Incidence	1:153
Variety	

1. Pre-eclampsia	76.28	%
2. Eclampsia	23.72	%
a. Antepartum	19.03	%
b. Postpartum	4.69	%

Parity:

Primi = 1:1.4 = Multigravidarum

Maternal mortality rate	3.85	%
(7 cases undelivered, 1 B.B.A.)		
Fetal mortality rate	9.25	%
neonatal death	1.28	%
intrauterine death	7.96	%

Placenta previa

Incidence	1:206	
Primigravidarum	11.05	%
Multigravidarum	88.93	%
Variety		
- total	39.9	%
- partial	38.3	%
- low insertion	22.2	%

Maternal mortality rate	2.54	%
Fetal mortality rate	40.45	%

Accidental haemorrhage

Incidence	1:503	
Primigravidarum	6.9	%
Multigravidarum	92.5	%
Maternal mortality rate	2.1	%
Fetal mortality rate	68.1	%

Ruptured uteri

Incidence	1:1,198	
Primigravidarum	10.4	%
Multigravidarum	98.7	%

Causes:

A. Spontaneous rupture

a. previous section scar	17.45	%
b. other causes	72.05	%

B. Obstetrical trauma

Maternal mortality rate	14.42	%
Fetal mortality rate	85.92	%

Maternal mortality rate
(in 1,000 live birth)


1. Due to obstetrical causes = 0.52:1,000 live birth
2. Due to associated diseases = 0.39:1,000 live birth
3. Due to toxic condition before admission = 0.34:1,000 live birth

Fetal mortality rate

In delivery room	= 20.21:1,000 live birth
1. Intrauterine death	15.6 : 1,000 live birth
2. Neonatal death	
- full term	2.59:1,000 live birth
- premature	2.02:1,000 live birth

STATISTICAL SUMMARY OF OBSTETRICAL DEPARTMENT
IN WOMEN'S HOSPITAL

The total number of deliveries (2494 - 2503)



<u>Multiple pregnancy</u>		<u>Method of deliveries</u>	
Twin	1:109	Normal deliveries	91.20 %
Triplet	1:8,724	Abnormal deliveries	8.80 %
Quadruplet	1:34,069	B.B.A. (305 cases)	0.31 %
Siamese twin	1:34,069		

<u>Sex</u>		<u>The total number of live birth</u>	
Male	52.26 %	Full term	77,378 = 95.79 %
Female	47.74 %	Premature	3,047 = 4.21 %

<u>Weight</u>	<u>Length</u>	<u>Congenital anomalies</u>	
3,102 gm.	49.77 cm.	1. Abnormality of extremities	37.9 %
<u>Weight of placenta</u>	<u>length of umbilicus</u>	2. " " of G.I. tract	26.1 %
579 gm.	49.77 cm.	3. " " of nervous system	20.3 %
<u>Fetal skull</u>		4. " " of urogenital system	5.3 %
<u>Diameter</u>	<u>Circumference</u>	5. " " of E.E.N.T.	2.1 %
M.O. 13.5 cm.	37.5 cm.	6. Miscellaneous	8.3 %
F.O. 11.0 "	34.0 "		

				<u>Position and presentation</u>	
S.O.B.	9.5 "	31.5 "		O.L.A.	62.9 %
B.P.	9.2 "			O.R.A.	33.81 %
B.T.	7.0 "			O.P.P.	0.80 %
				O.L.P.	0.43 %
				O.R.P.	0.18 %
				Breech	1.7 %
				Transverse	0.19 %
				Face	0.08 %
				Brow	0.0001 %

<u>Blood Kahn</u>			
	2494	2498	2502 2503
3+	-	-	0.62% 0.86%
4+	7.8%	7.18%	1.5 % 2.47%

Operative obstetric deliveries

Forceps extraction	2.49	%
Caesarean section	2.08	%
Craniotomy	0.04	%
Evisceration and Embryotomy	0.01	%
Internal version	0.23	%
Manual removal of placenta	0.70	%

Placenta previa

Incidence	1:206	
Primigravidarum	11.05	%
Multigravidarum	88.93	%
Variety		
- total	39.9	%
- partial	38.3	%
- low insertion	22.2	%

Postpartum haemorrhage

Incidence	2.05	
Grade I (300-500)	56.4	%
Grade II (500-1000)	37.4	%
Grade III (1000-1500)	5.8	%
Grade IV (1500-2000)	0.4	%

Maternal mortality rate	2.54	%
Fetal mortality rate	40.45	%

Accidental haemorrhageThe Toxemia of Pregnancy

Incidence	1:153	
Variety		
1. Pre-eclampsia	76.28	%
2. Eclampsia	23.72	%
a. Antepartum	19.03	%
b. Postpartum	4.69	%

Incidence	1:503	
Primigravidarum	6.9	%
Multigravidarum	92.5	%
Maternal mortality rate	2.1	%
Fetal mortality rate	68.1	%

Ruptured uteri

Parity:

Primi = 1:1.4 = Multigravidarum

Maternal mortality rate	3.85	%
(7 cases undelivered, 1 B.B.A.)		
Fetal mortality rate	9.25	%
neonatal death	1.28	%
intrauterine death	7.96	%

Causes:

A. Spontaneous rupture

a. previous section scar	17.45	%
b. other causes	72.05	%

B. Obstetrical trauma

Maternal mortality rate	14.42	%
Fetal mortality rate	85.92	%

Maternal mortality rate
(in 1,000 live birth)


1. Due to obstetrical causes	=	0.52:1,000 live birth
2. Due to associated diseases	=	0.39:1,000 live birth
3. Due to toxic condition before admission	=	0.34:1,000 live birth

Fetal mortality rate

In delivery room	=	20.21:1,000 live birth
1. Intrauterine death		15.6 : 1,000 live birth
2. Neonatal death		
- full term		2.59:1,000 live birth
- premature		2.02:1,000 live birth

STATISTICAL SUMMARY OF OBSTETRICAL DEPARTMENT
IN WOMEN'S HOSPITAL

The total number of deliveries (2494 - 2503)



<u>Multiple pregnancy</u>		<u>Method of deliveries</u>	
Twin	1:109	Normal deliveries	91.20 %
Tripet	1:8,724	Abnormal deliveries	8.80 %
Quadrupet	1:34,069	B.B.A. (305 cases)	0.31 %
Siamese twin	1:34,069		

<u>Sex</u>		<u>The total number of live birth</u>	
Male	52.26 %	Full term	77,378 = 95.79 %
Female	47.74 %	Premature	3,047 = 4.21 %

<u>Weight</u>	<u>Length</u>	<u>Congenital anomalies</u>	
3,102 gm.	49.77 cm.	1. Abnormality of extremities	37.9 %
<u>Weight of placenta</u>	<u>length of umbilicus</u>	2. " " of G.I. tract	26.1 %
579 gm.	49.77 cm.	3. " " of nervous system	20.3 %
<u>Fetal skull</u>		4. " " of urogenital system	5.3 %
<u>Diameter</u>	<u>Circumference</u>	5. " " of E.E.N.T.	2.1 %
M.O. 13.5 cm.	37.5 cm.	6. Miscellaneous	8.3 %
F.O. 11.0 "	34.0 "		
S.O.B. 9.5 "	31.5 "		
B.P. 9.2 "			
B.T. 7.0 "			

				<u>Position and presentation</u>	
				O.L.A.	62.9 %
				O.R.A.	33.81 %
				O.P.P.	0.80 %
				O.L.P.	0.43 %
				O.R.P.	0.18 %
				Breech	1.7 %
				Transverse	0.18 %
				Face	0.08 %
				Brow	0.0001 %

<u>Blood Kahn</u>				
	2494	2498	2502	2503
3+	-	-	0.62%	0.86%
4+	7.8%	7.18%	1.5 %	2.47%

Operative obstetric deliveries

Forceps extraction	2.49	%
Caesarean section	2.08	%
Craniotomy	0.04	%
Evisceration and Embryotomy	0.01	%
Internal version	0.23	%
Manual removal of placenta	0.70	%

Postpartum haemorrhage

Incidence	2.05	
Grade I (300-500)	56.4	%
Grade II (500-1000)	37.4	%
Grade III (1000-1500)	5.8	%
Grade IV (1500-2000)	0.4	%

The Toxemia of Pregnancy

Incidence	1:153	
Variety		
1. Pre-eclampsia	76.28	%
2. Eclampsia	23.72	%
a. Antepartum	19.03	%
b. Postpartum	4.69	%

Parity:

Primi = 1:1.4 = Multigravidarum

Maternal mortality rate	3.85	%
(7 cases undelivered, 1 B.B.A.)		
Fetal mortality rate	9.25	%
neonatal death	1.28	%
intrauterine death	7.96	%

Placenta previa

Incidence	1:206	
Primigravidarum	11.05	%
Multigravidarum	88.93	%
Variety		
- total	39.9	%
- partial	38.3	%
- low insertion	22.2	%

Maternal mortality rate	2.54	%
Fetal mortality rate	40.45	%

Accidental haemorrhage

Incidence	1:503	
Primigravidarum	6.9	%
Multigravidarum	92.5	%
Maternal mortality rate	2.1	%
Fetal mortality rate	68.1	%

Ruptured uteri

Incidence	1:1,198	
Primigravidarum	10.4	%
Multigravidarum	98.7	%

Causes:

A. Spontaneous rupture

a. previous section scar	17.45	%
b. other causes	72.05	%

B. Obstetrical trauma

Maternal mortality rate	14.42	%
Fetal mortality rate	85.92	%

Maternal mortality rate
(in 1,000 live birth)


1. Due to obstetrical causes = 0.52:1,000 live birth
2. Due to associated diseases = 0.39:1,000 live birth
3. Due to toxic condition before admission = 0.34:1,000 live birth

Fetal mortality rate

In delivery room	= 20.21:1,000 live birth
1. Intrauterine death	15.6 : 1,000 live birth
2. Neonatal death	
- full term	2.59:1,000 live birth
- premature	2.02:1,000 live birth

STATISTICAL SUMMARY OF OBSTETRICAL DEPARTMENT IN WOMEN'S HOSPITAL

The total number of deliveries (2494 - 2503)



<u>Multiple pregnancy</u>		<u>Method of deliveries</u>	
Twin	1:109	Normal deliveries	91.20 %
Tripet	1:8,724	Abnormal deliveries	8.80 %
Quadrupet	1:34,069	B.B.A. (305 cases)	0.31 %
Siamese twin	1:34,069		

<u>Sex</u>		<u>The total number of live birth</u>	
Male	52.26 %	Full term	77,378 = 95.79 %
Female	47.74 %	Premature	3,047 = 4.21 %

<u>Weight</u>	<u>Length</u>	<u>Congenital anomalies</u>	
3,102 gm.	49.77 cm.	1. Abnormality of extremities	37.9 %
<u>Weight of placenta</u>	<u>length of umbilicus</u>	2. " " of G.I. tract	26.1 %
579 gm.	49.77 cm.	3. " " of nervous system	20.3 %
<u>Fetal skull</u>		4. " " of urogenital system	5.3 %
<u>Diameter</u>	<u>Circumference</u>	5. " " of E.E.N.T.	2.1 %
M.O. 13.5 cm.	37.5 cm.	6. Miscellaneous	8.3 %

		<u>Position and presentation</u>	
F.O. 11.0 "	34.0 "	O.L.A.	62.9 %
S.O.B. 9.5 "	31.5 "	O.R.A.	33.81 %
B.P. 9.2 "		O.P.P.	0.80 %
B.T. 7.0 "		O.L.P.	0.43 %
		O.R.P.	0.18 %
		Breech	1.7 %
		Transverse	0.18 %
		Face	0.0001 %
		Brow	

<u>Blood Kahn</u>			
	2494	2498	2502 2503
3+	-	-	0.62% 0.36%
4+	7.8%	7.18%	1.5 % 2.47%

Operative obstetric deliveries

Forceps extraction	2.49	%
Caesarean section	2.08	%
Craniotomy	0.04	%
Evisceration and Embryotomy	0.01	%
Internal version	0.23	%
Manual removal of placenta	0.70	%

Postpartum haemorrhage

Incidence	2.05	
Grade I (300-500)	56.4	%
Grade II (500-1000)	37.4	%
Grade III (1000-1500)	5.8	%
Grade IV (1500-2000)	0.4	%

The Toxemia of Pregnancy

Incidence	1:153	
Variety		

1. Pre-eclampsia	76.28	%
2. Eclampsia	23.72	%
a. Antepartum	19.03	%
b. Postpartum	4.69	%

Parity:

Primi = 1:1.4 = Multigravidarum

Maternal mortality rate	3.85	%
(7 cases undelivered, 1 B.B.A.)		
Fetal mortality rate	9.25	%
neonatal death	1.28	%
intrauterine death	7.96	%

Placenta previa

Incidence	1:206	
Primigravidarum	11.05	%
Multigravidarum	88.93	%
Variety		
- total	39.9	%
- partial	38.3	%
- low insertion	22.2	%

Maternal mortality rate	2.54	%
Fetal mortality rate	40.45	%

Accidental haemorrhage

Incidence	1:503	
Primigravidarum	6.9	%
Multigravidarum	92.5	%
Maternal mortality rate	2.1	%
Fetal mortality rate	68.1	%

Ruptured uteri

Incidence	1:1,198	
Primigravidarum	10.4	%
Multigravidarum	98.7	%

Causes:

A. Spontaneous rupture

a. previous section scar	17.45	%
b. other causes	72.05	%

B. Obstetrical trauma

Maternal mortality rate	14.42	%
Fetal mortality rate	85.92	%

Maternal mortality rate
(in 1,000 live birth)

1. Due to obstetrical causes = 0.52:1,000 live birth
2. Due to associated diseases = 0.39:1,000 live birth
3. Due to toxic condition before admission = 0.34:1,000 live birth

Fetal mortality rate

In delivery room	= 20.21:1,000 live birth
1. Intrauterine death	15.6 : 1,000 live birth
2. Neonatal death	
- full term	2.59:1,000 live birth
- premature	2.02:1,000 live birth

เอกสารส่วนบุคคล ศาสตราจารย์นายแพทย์เสม พริ้งพวงแก้ว

สบบ 1

.2/1.9 สติศึกษาจากผลงานศิลปะ แผนกจิตรกรรม (ชื่อเรื่อง)
พ.ศ. 2497 โรงพยาบาลเสนาบดี

พ.ศ. 2497

จำนวน 13 แผ่น

แฟ้มที่ 2


กล่องที่ 1

STATISTICS 2497

OBSTETRICAL DEPARTMENT

AVERAGE VALUES

Pelvis	23.5	26	18 cms.
Placenta	602.7		gms.
Fetus	3152		"
C.H.L.	50.5		cms.
Umbilical cord	49.9		"




มาตราส่วน 15 : 100

แสดงสถิติการคลอดผิดปกติ ปี พ.ศ. ๒๕๐๑

(รวม Abruptio placenta)

จำนวนทั้งหมด ๕๐๖

(Dead fetus มี ๘๖ คิดเป็น: dead เกิดจาก abnormal deliveries เท่านั้น)



มกราคม กุมภาพันธ์ มีนาคม เมษายน พฤษภาคม มิถุนายน กรกฎาคม สิงหาคม กันยายน ตุลาคม พฤศจิกายน ธันวาคม


แสดงจำนวนเปรียบเทียบระหว่างคลอดปกติและผิดปกติของแต่ละเดือน

มาตราส่วน 1: 500

ในปี พ.ศ. 2497

คลอดปกติ 544 คน

คลอดผิดปกติ 406 คน




Male Female Primigravida Multipara Abn. la. Prema. Kahn positive
52.25% 47.75% 34.76% 65.24% 7.63% 5.6% 7.54%


มาตราส่วน 1: 400

(จำนวนทั้งหมด)

สถิติเกิดคลอดปี พ.ศ. 2497



* Multiple pregnancy * P. P. H. * Uteri rupture * Tox of pregnancy * Placenta previa * Breech * O. P. P. * Forceps extractio * Caesarian section * Internal version * Airuptio placenta *



2495
2496
2497

เปรียบเทียบการคลอดผิดปกติ ในระหว่าง 3 ปี

2495 - 2496 - 2497

STATISTICS 2497
OBSTETRIC DEPARTMENT

Total numbers	6158	Cases
Normal deliveries	5447	"
Premature deliveries	305	"
Abnormal deliveries	406	"

NORMAL DELIVERIES

January	429	Cases
February	418	"
March	416	"
April	427	"
May	459	"
June	459	"
July	431	"
August	451	"
September	458	"
October	543	"
November	451	"
December	505	"

Primigravida	34.76	%
Multigravida	65.24	%
Male	52.25	%
Female	47.75	%

O.L.A.	74.6	%
O.R.A.	25.4	%

Average Values

Weight	3152.15	gms.
C.H.L.	50.51	cms.
Placenta	602.75	gms.
Umbilical cord	49.98	cms.

External pelvic measurment 23.533, 25.596, 18.067
23.5 , 26 , 19

Heaviest weight	5100	gms.
C.H.L. Longest	54	cms.

Shortest	45	cms.
Placenta :- Lightest	282	gms.

Heaviest	1050	gms.
Umbilical cord :- Longest	103	cms.

Shortest	11	cms.
----------	----	------

Blood Kahn

Kahn negative	84.79	%
Kahn positive	7.54	%
Unknown	7.67	%

ABNORMAL DELIVERIES2497MULTIPLE PREGNANCIES

<u>Total cases</u>	55	= 1.1 %
Twins	52	
Triplets	1	
Quaduplets ^r	1	
Siamese twins(Thor ^a copagus) ₊	1	

DEAD FETUS

<u>Total cases</u>	141	= 2.82 %
Maceration		
Full term	26	
Premature	17	
Still born	7	
Asphyxia	3	
Dead due to abnormal deliveries	86	
Dead in ward	2	

POSTPARTUM HEMORRHAGE

<u>Total cases</u>	54	= 1.08 %
Primigravida	10	
Multigravida	44	
Causes :-		
Retained placenta	14	
Uteri inertia	22	
Unknown	2	
Delivery at home	16	
Maternal mortality	4	= 7.4 %

PLACENTA PREVIA

<u>Total cases</u>	27	= 0.54 %
Primigravida	4	
Multigravida	23	
<u>Managements</u>		
Caesarean section	24	
Forceps extraction	0	
Normal labour	3	
Fetal mortality	17	
Still born	2	
Maceration	15	
Maternal mortality	1	= 3.7 %
(due to psychosis P.O.)		

ABNORMAL POSITION

<u>Total cases</u>	120	= 2.4 %
Primigravida	46	
Multigravida	74	
Breech presentation	90	= 1.8 %
S.R.A.	35	
S.L.A.	55	
Occiput posterior	29	= 0.58%
Face presentation	1	= 0.02%
Fetal mortality	7	
Still born	4	
(Hydrocephalus 1, Breech 2)		
Maceration	3	

FORCEPS EXTRACTION

<u>Total cases</u>	77	= 1.54 %
Primigravida	57	
Multigravida	20	
<u>Indications</u>		
Prolong second stage	60	
Uteri inertia	6	
Eclampsia	8	
Prolapse cord	1	
Congenital heart disease	1	
Twins (second baby)	1	

Fetal mortality

Still born 4

Maceration 2

Maternal mortality 1 = 1.3 %

(due to P.P.H.)

UTERI RUPTURETotal cases 2 = 0.04 %

Primigravida 0

Multigravida 2

Causes

Old scar of Caesarean section 1

Transverse presentation 1

Fetal mortality 1Maternal mortality 1 = 50 %TOXEMIA OF PREGNANCYTotal cases 37 = 0.74 %

Primigravida 24

Multigravida 13

Types

Eclampsia 20

Hydreds 16

Postpartum eclampsia 1

Fetal mortality 6Maternal mortality 4 = 10.81%Managements

Caesarean section 21

Forceps extraction 8

Normal labour 5

Symptomtics 3

CAESAREAN SECTION

<u>Total cases</u>	112	= 2.24 %
Primigravida	50	
Multigravida	62	
<u>Types</u>		
Classical	101	
Low cervical	5	
Low transverse	6	
<u>Indications</u>		
Placenta previa	24	
Prolong ^{ed} second stage	23	
Toxemia of pregnancy	21	
Repeated Caesarean section	9	
Transverse presentation	6	
Elderly primigravida	6	
Abruptio placenta	6	
Uteri inertia	4	
Twins	3	
Prolong ^{ed} first stage	3	
General contracted pelvis	2	
Over term pregnancy	2	
Face presentation	1	
Epilepsy	1	
Missed abortion	1	

Fetal mortality 20

Maternal mortality 3 = 2.67%

(due to abruptio placenta 1,
placenta previa 1,uteri inertia P.O. 1)

INTERNAL VERSION

<u>Total cases</u>	9	=	0.18 %
Primigravida	1		
Multigravida	8		
<u>Indications</u>			
Prolapse arm	4		
Transverse presentation	3		
Prolapse cord	1		
Twins (second baby)	1		
<u>Fetal mortality</u>	8		
<u>Maternal mortality</u>	1	=	11.11%
(report in P.P.H.)			

ABRUPTIO PLACENTA

<u>Total cases</u>	6	=	0.12%
Primigravida	1 (Twins)		
Multigravida	5		
<u>Fetal mortality</u>	4		
<u>Maternal mortality</u>	1	=	16.67%

ANORMALIESTotal cases

34

Stiffness of neck	2
Small nodule at rt. breast	1
Intrauterine amputation - both hands	2
Small nodule at rt. big toe	1
Polydactylism	1
Enlargement of rt. calf	1
Phimosis	1
Anencephalus	4
Small nodule at left foot	1
Congenital megacolon	1
Hydrocephalus	3
Nodule at scalp	1
Telepes	3
Harelip	3
Brachial paralysis	1
Angulation of both wrist joints	1
Facial paralysis	1
Hydrocele	1
Imperforated anus	1
Rectovaginal fistula	1
Nodule at right ear	1
Flat foot	1
Shortness of neck	1

COMPLICATIONSTotal cases

49

Ber-beri	5
Bronchitis	3
Pharyngitis	1
Hepatitis	1
Lobar pneumonia	2
Puerperal fever	11
Schizophrenia	1
Hypertension	1
Cerebral malaria	1
Rheumatic arthritis	1
Pulmonary infiltration cause?	1
Poly-arthritis	1
Mitral stenosis	1
Isolated pulmonary stenosis	1
Anemia	4
Pleural effusion	1
Malaria	2
Root abscess	1
Cardiac arrhythmia th	1
Colitis	1
Condyloma acuminata	1
Cracked nipple	1
Chicken-pox	2
Pulm.T.B.	2
Rheumatic heart	1
Hematoma of rt. labia	1

เอกสารส่วนบุคคล ศาสตราจารย์นายแพทย์เสมอ พริ้งพวงแก้ว

ឥប 1

2/1.10 สภากาชาดไทย ถนนวิภาวดี 2496 (ชื่อเรื่อง)
-2499 โรงพยาบาลสงฆ์และสัตว์

พ.ศ. ๒๕๐๑

จำนวนแผ่น

แฟ้มที่ 2

กล่องที่ 1

175001
Beriberi-Pregnancy.

W. H. 1882

6 100

W. H. 1883

6 100

W. H. 1884

7 100

W. H. 1885

—

(ส.ว.)

ที่ ๗๗ / ๒๕๐๑

โรงพยาบาลหญิง พญาไท

๑๙ มกราคม ๒๕๐๑

เรื่อง จำนวนผู้ป่วยเป็นโรคเหน็บชา

เรียน ผู้อำนวยการกองส่งเสริมอาหาร

อ้างถึง หนังสือที่ ๑๔๘๓/๒๕๐๐ ลงวันที่ ๓๐ พฤศจิกายน ๒๕๐๐

สิ่งที่ส่งมาด้วย สถิติผู้ป่วยเป็นโรคเหน็บชา โรงพยาบาลหญิง ๑ ฉบับ
โรงพยาบาลเด็ก ๑ ฉบับ

ตามหนังสือที่อ้างถึง กองส่งเสริมอาหารมีความประสงค์ใคร่จะทราบ

จำนวนผู้ป่วยเป็นโรคเหน็บชา ซึ่งมารับการบำบัดรักษาในโรงพยาบาลแห่งนี้ เพื่อประกอบการพิจารณาหาทางป้องกันให้ร้กมยิ่งขึ้น นั้น

บัดนี้ เจ้าหน้าที่สถิติของโรงพยาบาลได้รวบรวมผู้ป่วยเป็นโรคเหน็บชา โดยแยกประเภทคนไข้นอก คนไข้ใน อายุ และเพศ ตามที่กองส่งเสริมอาหารต้องการ เรียบมาเพื่อทราบพร้อมหนังสือนี้ด้วยแล้ว.

ขอแสดงความนับถืออย่างสูง

(๑๕๕๐) ส.ว.

(นายแพทย์เสม พริ้งพวงแก้ว)

นายแพทย์ผู้อำนวยการโรงพยาบาลหญิง

รับทราบ ๑๕/๑/๕๐๑



ส.ว. ๑๕/๑/๕๐๑

๓ ก.พ. ๒๕๐๑

สถิติคนไข้ที่เป็นโรคเหน็บชา. ซึ่งมารับการบำบัดรักษาในโรงพยาบาลหญิง
ปี ๒๔๔๔

จำนวนคนไข้ภายใน ๒๓ คน

อายุ	จำนวน
๑๖ ปี	๑ คน
๑๘ "	๒ "
๒๑ "	๓ "
๒๒ "	๑ "
๒๓ "	๑ "
๒๔ "	๑ "
๒๕ "	๑ "
๒๖ "	๑ "
๒๗ "	๑ "
๒๘ "	๑ "
๓๗ "	๒ "
๓๙ "	๑ "
๔๐ "	๒ "
๔๑ "	๒ "
๔๒ "	๒ "
๕๐ "	๑ "
๕๐ "	๒ "
จำนวนคนไข้ภายนอก	๑๔๔ คน

สถิติคนไข้ที่เป็นโรคเหน็บชา ซึ่งมารับการบำบัดรักษาใน ร.พ. เด็ก ปี ๒๔๔๔ จำนวนคนไข้ภายใน.

อายุ.	เพศ.	จำนวน.
๑๐ วัน	ชาย	๑ คน
๒ เดือน	ชาย	๑ คน
๓ เดือน	ชาย	๔ คน
๑ ปี	ชาย	๑ คน
๑ เดือน	หญิง	๑ คน
๓ ปี	หญิง	๑ คน

จำนวนคนไข้ภายนอก.

ชาย ๒๓ คน

หญิง ๒๗ คน.